DOCUMENT RESUME

ED 288 336 EC 201 217

AUTHOR Patrias, Karen

TITLE Newborn Screening: National Library of Medicine

Literature Search, January 1980 through March 1987.

No. 87-2.

INSTITUTION National Library of Medicine (DHHS/NIH), Bethesda,

Md.

PUB DATE Mar 87 NOTE 58p.

AVAILABLE FROM Literature Search Program, Reference Section,

National Library of Medicine, 8600 Rockville Pike,

Bethesda, MD 20894 (free; send a gummed,

self-addressed label).

PUB TYPE Reference Materials - Bibliographies (131)

EDRS PRICE MF01/PC03 Plus Postage.

DESCRIPTORS *Congenital Impairments; *Disabilities; *Handicap

Identification; Metabolism; *Neonates; *Screening Tests; Sickle Cell Anemia; Special Health Problems

IDENTIFIERS Cystic Fibrosis; Galactosemia; Hypothyroidism;

Muscular Dystrophy; Phenylketonuria

ABSTRACT

This bibliography, prepared by the National Library of Medicine through a literature search of its online databases, covers all aspects of newborn screening. It includes references to screening for: inborn errors of metabolism, such as phenylketonuria and galactosemia; hemoglobinopathies, particularly sickle cell disease; congenital hypothyroidism and congenital adrenal hyperplasia; cystic fibrosis; and muscular dystrophy. Citations are primarily to journal articles from "MEDLINE," "HEALTH PLANNING AND ADMINISTRATION," and "BIOETHICSLINE." The bibliography contains over 750 references to English and foreign-language publications, issued between January 1980 and March 1987, in alphabetic order. (Author/JDD)

Reproductions supplied by EDRS are the best that can be made from the original document.

NATIONAL LIBRARY of MEDICINE

No. 87-2

LITERATURE SEARCH

NEWBORN SCREENING

January 1980 through March 1987

753 Citations

Prepared by Karen Patrias

U.S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it.
- Minor changes have been made to improve reproduction quality
- Points of view or opinions stated in this document do not necessarily represent official OERI position or policy

Screening of infants in the first month of life for inherited metabolic disorders and other diseases requiring early diagnosis has been known for over thirty years, beginning with programs in the 1950's for detection of phenylketonuria (PKU). Although many advances have been made both in the numbers of disorders for which screening tests have been developed and in the sophistication and accuracy of the tests themselves, controversy currently exists over what constitutes an ideal newborn screening program. Specific issues include technical as well as socioeconomic questions such as what diseases should be included, which are the most effective testing methods, should screening programs be voluntary or mandatory and for what groups of infants, and what is their cost-benefit to society as a whole.

This bibliography covers all aspects of newborn screening and includes specific reference to screening for inborn errors of metabolism, such as PKU and galactosemia; hemoglobinopathies, particularly sickle cell disease; congenital hypothyroidism and congenital adrenal hyperplasia; cystic fibrosis; and muscular dystrophy. Citations are primarily to journal articles from NLM's MEDLINE, HEALTH PLANNING AND ADMINISTRATION, and BIOETHICSLINE databases.

"Newborn Screening" was prepared in part in support of the NIH Consensus Development Conference on Newborn Screening for Sickle Cell Disease and Other Hemoglobinopathies, held in Bethesda, MD, April 6-8, 1987.

The publications of the National Library of Medicine's (NLM) Literature Search series contain citations to journal articles and monographs selected from a variety of online databases. NLM databases utilized include MEDLINE, BIOETHICSLINE, CANCERLIT, HEALTH PLANNING AND ADMINISTRATION, TOXLINE, and CATLINE, among others.

Citations are formatted according to the rules established for publication of *Index Medicus*. A full explanation of the elements as they are included in the citations may be found in each issue of that publication as well as in *Abridged Index Medicus*. A bracketed title indicates that the article is in a foreign language. For these foreign titles, the language of publication is shown by a three letter abbreviation at the end of the citation.

This publication is not copyrighted and may be freely reproduced. A complete list of all available Literature Searches appears in each issue of *Index Medicus* and *Abridged Index Medicus*. For a single copy of any of the other publications in this series, send a gummed, self-addressed label to: Literature Search Program, Reference Section, National Library of Medicine, 8600 Rockville Pike, Bethesda, MD 20894.

SEARCH STRATEGY

The formulation used for searching MEDLINE for this Literature Search is given below. Please note that search strategies for the Literature Searches differ from individual demand searches in that they are generally broadly formulated and irrelevant citations edited out prior to printing.

- SS 1 = INFANT, NEWBORN OR INFANT, NEWBORN, DISEASES/DIAGNOSIS
- SS 2 = 1 AND MASS SCREENING OR 1 AND DIAGNOSTIC TESTS, ROUTINE
- SS 3 = 1 AND ALL SCREEN:
- SS 4 = EXP METABOLISM, INBORN ERRORS
- SS 5 = EXP HYPOTHYROIDISM OR ADRENAL HYPERPLASIA, CONGENITAL
- SS 6 = EXP HEMOGLOBINOPATHIES OR IGG OR ALL BIOTIN:
- SS 7 = CYSTIC FIBROSIS OR MUSCULAR DYSTROPHY
- SS 8 = 3 AND 4 OR 3 AND 5 OR 3 AND 6 OR 3 AND 7
- SS 9 = (TW) ALL NEWBORN: OR ALL NEONAT:
- SS10 = 9 AND ALL SCREEN: (TW)
- SS11 = TS (TI) :SCREEN:
- SS12 = 2 OR 8 OR 11
- SS13 = 12 AND NOT ANIMAL

A

- Abuelo D; Pueschel S; Stern L. Screening the neonate for development and genetic disorders. Paediatrician 1982;11(1-2):45-71
- Acosta PB. Prevention of mental retardation through screening, retrieval, diagnosis, and management of inherited metabolic disease. Ala J Med Sci 1982 Oct;19(4 Suppl 1):10-6
- Adelman CS. The constitutionality of mandatory genetic screening statutes. Case West Reserve Law Rev 1981 Summer;31(4):897-948
- Adriaenssens K; Janssens H; van Soom H. Two tier screen for cystic fibrosis [letter]. Lancet 1981 Apr 11;1(8224):833
- Adriaenssens K; Van Riel L. Serum pancreatic lipase as a screening test for cystic fibrosis. Arch Dis Child 1982 Jul;57(7):553-5
- Adriaenssens K; Vermeiren G. Simple electrophoretic technique for creatine kinase MM isozyme in neonatal Duchenne muscular disease screening using dried blood samples. Clin Chim Acta 1980 Jul 17;105(1):99-103
- Adriaenssens KM; Philips ES; Colfs B. A quantitative IgE radioimmunoassay in dried blood spots suitable for neonatal screening of atopy. Clin Chim Acta 1985;151(1):91-5
- Ahmed P; Ahmad KN. Screening of the newborns for glucose-6-phosphate dehydrogenase deficiency. Indian Pediatr 1983;20(5):351-5
- Al Awamy BH; Al Muzan M; Al Turki M; Serjeant GR. Neonatal screening for sickle cell disease in the Eastern Province of Saudi Arabia. Trans R Soc Trop Med Hyg 1984;78(6):792-4
- Alberti PW; Hyde ML; Riko K; Corbin H; Abramovich S. An evaluation of BERA for hearing screening in high-risk neonates. Laryngoscope 1983 Sep;93(9):1115-21
- Albertini A; Bugiardini G; Carta S; Giovannelli G; Grandolfo ME; Pinchera A; Sereni Piceni L; Sorcini M; Tomarchio S; Varrone S. External quality control for neonatal thyroid screening [letter]. J Endocrinol Invest 1985 Oct;8(5):485-6
- Allan WC; Philip AG. Neonatal cerebral pathology diagnosed by ultrasound. Clin Perinatol 1985 Feb;12(1):195-218
- Allhoff P; Weidtman V. [First examination of neonates: Date of an early disease diagnosis programme]. Dtsch Med Wochenschr 1985;110(3):91-6 [Ger]
- Alm J; Hagenfeldt L; Larsson A; Lundberg K. Incidence of congenital hypothyroidism: retrospective study of neonatal laboratory screening versus clinical symptoms as indicators leading to diagnosis. Br Med J [Clin Res] 1984 Nov 3;289(6453):1171-5

- Alm J; Holmgren G; Larsson A; Schimpfessel L. Histidinaemia in Sweden.
 Report on a neonatal screening programme. Clin Genet 1981;20(3):229-33
- Alm J; Larsson A. Evaluation of a nation-wide neonatal metabolic screening programme in Sweden 1965-1979. Acta Paediatr Scand 1981;70(5):601-7
- Alm J; Larsson A; Rosenqvist U. Health economic analysis of the Swedish neonatal metabolic screening programme. A method of optimizing routines. Med Decis Making 1982;2(1):33-45
- Alm J; Larsson A; Westphal O. [A program for the nation-wide screening of newborn infants for congenital hypothyreosis]. Lakartidningen 1980 Aug 27;77(35):2884-6 [Swe]
- Amanullah A; Hanash S; Bunnell K; Strahler J; Rucknagel DL; Ferruci SJ. Cord blood screening for hemoglobin disorders by high-performance liquid chromatography. Anal Biochem 1982 Jul 1:123(2):402-7
- American Academy of Pediatrics Committee on Genetics: New issues in newborn screening for phenylketonuria and congenital hypothyroidism.

 Pediatrics 1982 Jan;69(1):104-6
- Anakura M. [Clinical studies of histidinemia (author's transl)] Hokkaido Igaku Zasshi 1981 Jan;56(1):1-15 [Jpn]
- Anders GJPA. [Neonatal screening in the Netherlands]. J Genet Hum 1981; 29(1):23-6 [Fre]
- Andrew M; Karpatkin M. A simple screening test for evaluating prolonged partial thromboplastin times in newborn infants. J Pediatr 1982 Oct; 101(4):610-2
- Andrews, Lori B. Legal Liability and Quality Assurance in Newborn Screening. Chicago: American Bar Foundation, 1985. 242 p.
- Andrews, Lori B. State Laws and Regulations Governing Newborn Screening. Chicago: American Bar Foundation, 1985. 167 p.
- Anionwu E. Sickle cell disease. Health Visit 1982 Jul;55(7):336-41
- Anionwu EN. Sickle cell disease: screening and counselling in the antenatal and neonatal period. Part 2. Midwife Health Visit Community Nurse 1983 Nov;19(11):440-3
- Anionwu EN. Sickle cell disease: screening and counselling in the antenatal and neonatal period. Part I. Midwife Health Visit Community Nurse 1983 Oct;19(10):402-6
- Annas GJ. Mandatory PKU screening: the other side of the looking glass. Am J Public Health 1982 Dec;72(12):1401-1403

- Antonozzi I; Dominici R; Andreoli M; Monaco F. Neonatal screening in Italy for congenital hypothyroidism and metabolic disorders:

 Hyperphenylalaninemia, maple syrup urine disease and homocystinuria. J Endocrinol Invest 1980;3(4):357-63
- Appleton RE; Baumer JH; Holton JB. A loophole in the phenylketonuria screening programme [letter]. Lancet 1984 Sep 29;2(8405):752
- Arends J; Nrgaard, Pedersen B. Immunofluorometry of thyrotropin, from whole-blood spots on filter paper, to screen for congenital hypothyroidism. Clin Chem 1986 Oct;32(10):1854-6
- Arnold MB; Arulanantham K; Bapat V; et al. Characteristics of infantile hypothyroidism discovered on neonatal screening. J Pediatr 1984; 104(4):539-44
- Arnold MB; Arulanantham K; Bapat V; et al. Effects of neonatal screening for hypothyroidism: Prevention of mental retardation by treatment before clinical manifestations. Lancet 1981;2(8255):1095-8
- Arslan E; Prosser S; Conti G. [Clinical applications of evoked auditory potentials in pediatric audiology]. Acta Otorhinolaryngol Ital 1984;4 Suppl 3:101-10 [Ita]
- Asami T. Screening for hypercholesterolaemia on blood spotted filterpaper [letter]. Lancet 1983 Jul 23;2(8343):229-30
- Ashkenazi S; Mimouni F; Merlob P; Reisner SH. Neonatal biliruben levels and glucose-6-phosphate dehydrogenase deficiency in preterm and low-birth-weight infants in Israel. Isr J Med Sci 1983 Dec;19(12):1056-8
- Aula P. [Are hereditary diseases avoidable?] Duodecim 1986;102(16):1094-9 [Fin]
- Avni EF; Brion L; Daled HJ. Ultrasonic screening evaluation of intracranial pathology in "high risk" neonates. 100 cases experience. J Belge Radiol 1981;64(6):487-94

В

- Babin JP; Martin Cl. [Diagnosis and screening of congenital hypothyroidism in newborns: Elements of neonatal thyroid physiology]. Bordeaux Med 1980;13(1-3):3-8 [Fre]
- Baginski ES; Epstein E; Zak B. Some aspects of bilirubin determination in the newborn using dimethyl sulfoxide. Ann Clin Lab Sci 1980;10(6):486-92
- Baiget M; Del Rio E; Domenech M; et al. [Hemoglobinopathies screening in umbilica! cord blood]. Biol Clin Hematol 1981;3(4):251-6 [Spa]

- Baird AS; Hemming AM. Neonatal vision screening. J Visual Impair Blind 1982 May;76(5):182-5
- Barden HS; Kessel R. The costs and benefits of screening for congenital hypothyroidism in Wisconsin. Soc Biol 1984 Fall-Winter;31(3-4):185-200
- Barden HS; Kessel R; Schuett VE. The costs and benefits of screening for PKU in Wisconsin. Soc Biol 1984 Spring-Summer; 31(1-2):1-17
- Barnes ND. Screening for congenital hypothyroidism: the first decade. Arch Dis Child 1985 Jun;60(6):587-92
- Baumann P; Wood WG. Serum thyroglobulin concentrations in the first weeks of life as measured with an immunoluminometric assay. J Clin Chem Clin Biochem 1985 Nov;23(11):753-8
- Bednar J. [New information on methods of laboratory diagnosis of diseases of the thyroid gland. III. Screening for congenital hypothyroidism.

 Determination of thyrotropin]. Cesk Farm 1985 Mar;34(2):79-84 [Cze]
- Bellman S. Hearing screening in infancy [editorial]. Arch Dis Child 1986 Jul;61(7):637-8
- Belotti BM; Tarantino V; Tassone F. [Neonatal audiometric screening: A 5 year experience]. Gaslini 1982;14(1):55-9 [Ita]
- Bennett MJ; Bradley CE. Simpler liquid-chromatographic screening for organic acid disorders. Clin Chem 1984 Apr;30(4):542-6
- Berardi R; Baracchi MR; Borgogni P; Margollicci MA; Mattei R; Fois A. [Results of a screening project for congenital hypothyroidism in 4 years of experience]. Pediatr Med Chir 1982 Nov-Dec;4(6):657-60 [Ita]
- Berman BW; Ritchey AK; Jekel JF; Schwartz AD; Guiliotis DK; Pearson HA. Hematology of beta-thalassemia trait-age-related developmental aspects and intrafamilial correlations. J Pediatr 1980 Dec;97(6):901-5
- Berman JL. PKU-hypothesis concerning failures of screening [letter]. J Pediatr 1980 May;96(5):953-4
- Berman L; Klenerman L. Ultrasound screening for hip abnormalities: preliminary findings in 1001 neonates. Br Med J [Clin Res] 1986 Sep 20;293(6549):719-22
- Bernstein RE. Congenital hypothyroidism. Clinicopathological aspects and biochemical screening. S Afr Med J 1981;59(20):710-4
- Berry H K; Porter LJ. Newborn screening for phenylketonuria [letter].
 Pediatrics 1982 Sep;70(3):505-6
- Bhagwat GP; Mwisiya M. Glucose-6-phosphate dehydrogenase deficiency: screening of 200 Zambian newborn. Med J Zambia 1984 Jun;18(2):10-1

- Bhargava SK; Sachdev HPS; Iyer PU; Ramji S. Current status of infant growth measurements in the perinatal period in India. Acta Paediatr Scand 1985;74(Suppl. 319):103-10
- Bhattacharya J; Bennett MJ; Tucker SM. Long term follow up of newborns tested with the auditory response cradle. Arch Dis Child 1984 Jun; 59(6):504-11
- Bickel H. [Screening for congenital metabolic disorders. Indication and results]. Monatsschr Kinderheilkd 1983 Jun;131(6):323-7 [Ger]
- Bickel H; Bachmann C; Beckers R; et al. Neonatal mass screening for metabolic disorders. Summary of recent sessions of the Committee of Experts to Study Inborn Metabolic Diseases, Public Health Committee, Council of Europe. Eur J Pediatr 1981;137(2):133-9
- Bickel H; Schonberg D; Klett M. Neonatal thyroid screening [editorial].

 Eur J Pediatr 1984 Nov;143(1):6
- Blank RH. Public policy implications of human genetic technology: genetic screening. J Med Philos 1982 Nov;7(4):355-74
- Bliesener JA. [Ultrasonographic screening of the cranium in risk newborn (author's transl)]. Rontgenblatter 1980 Dec;33(12):626-31 [Ger]
- Boer HR. Hearing screening in "high risk" infants. NC Med J 1984 Mar; 45(3):157-8
- Bohles H; Stehr K. [Early diagnosis of congenital metabolic diseases]. Fortschr Med 1980 Jul 3;98(25):955-7 [Ger]
- Borgstrom A; Sveger T; Lindberg T; et al. Immunoreactive trypsin screening for cystic fibrosis. Acta Paediatr Scand 1982;71(4):621-4
- Boschetti R. [Neonatal screening (phenylketonuria and hypothyroidism) in France (1980)]. J Genet Hum 1981;29(1)27-9 [Fre]
- Bourdoux P; Branders C; Delange F; Ermans AM. Parenteral nutrition may interfere in thyrotropin screenings for congenital hypothyroidism [letter]. Clin Chem 1981 Nov;27(11):1943-4
- Bourguignon JP; Deby Dupont G; Reuter A; Senterre J; Lambotte C; Gerard A; Franchimont P. [Neonatal screening of mucoviscidosis using radioimmunologic levels of trypsin in dried blood cluates]. Rev Med Liege 1984 May 15;39(10):451-9 [Fre]
- Bowling FG; Brown AR. Development of a protocol for newborn screening for disorders of the galactose metabolic pathway. J Inherited Metab Dis 1986;9(1):99-104
- Bowling FG; Fraser DK; Clague AE; Hayes A; Morris DJ. A case of uridine diphosphate galactose-4-epimerase deficiency detected by neonatal screening for galactosaemia. Mcd J Aust 1986 Fcb 3;144(3):150-1

- Bozkowa K; Cabalska B; Duczynska N; Grodzka Z; Sendecka E; Nowakowska A; Lenartowska I; Kasperska-Dworak A; Helwich E. [Progress in the early detection of inborn errors of metabolism]. Probl Med Wieku Rozwoj 1981;10:69-85 [Pol]
- Brandt NJ; Jacobsen BB; Hummer L; et al. [Screening for congenital hypothyroidism in Denmark. II. Clinical findings in neonates diagnosed during a two-year period]. Ugeskr Laeg 1982;144(14):1004-7 [Dan]
- Briard ML. [Neonatal screening of phenylketonuria in France. Statistical data]. Arch Fr Pediatr 1983;40 Suppl 1:215-7 [Fre]
- Brusilow SW; Valle DL; Batshaw ML; Waber LJ. Screening for lethal genetic disease. Pediatrics 1982 Oct;70(4):647-8
- Buehler BA; Gortatouski MJ; Scow G; Hoogasian AC; Van Dyck PC. Thyroid screening in the newborn: Utah experience. Ann Clin Lab Sci 1983 Jan-Feb;13(1):5-9
- Busanny W; Grosse KP; Rziha HJ; Zur Hausen H. [Results of precise screening for prenatal cytomegalovirus and BK virus infections in newborn and young infants. II. BK virus infections]. Monatsschr Kinderheilkd 1980 May;128(5):269-70 [Ger]
- Butler R; Gitzelmann R; Illig R. [Screening of neonates for inherited metabolic diseases and congenital hypothyroidism in Switzerland].

 Arztl Lab 1981;27(5):119-23 [Ger]

Cacciari E; Balsamo A; Cassio A; et al. Neonatal screening programme for

- Cacciari E; Balsamo A; Cassio A; et al. Neonatal screening programme for congenital adrenal hyperplasia in a homogenous Caucasian population. J Inherit Metab Dis 1986;9(Suppl.1):142-6
- Cacciari E; Balsamo A; Cassio A; Piazzi S; Bernardi F; Salardi S;
 Cicognani A; Pirazzoli P; Zappulla F; Capelli M; et al. Neonatal
 screening program for congenital adrenal hyperplasia in a homogeneous
 Caucasian population. Ann NY Acad Sci 1985;458:85-9
- Cacciari E; Balsamo A; Cassio A; Piazzi S; Bernardi F; Salardi S; Cicognani A; Pirazzoli P; Zappulla F; Capelli M; et al. Neonatal screening for congenital adrenal hyperplasia. Arch Dis Child 1983 Oct;58(10):803-6
- Cacciari E; Balsamo A; Cassio A; Piazzi S; Bernardi F; Salardi S; Cicognani A; Pirazzoli P; Zappulla F; Capelli M; Paolini M.

 Neonatal screening for congenital adrenal hyperplasia using a microfilter paper method for 17-alpha-hydroxyprogesterone radioimmunoassay. Experience gained from the study of 22,233 cases. Horm Res 1982;16(1):4-9

- Cacciari E; Balsamo A; Piazzi S; Salardi S; Pirazili P; Capelli M;
 Cassio A; Bernardi F; Cicognani A; Zappulla F; Paolini M. Neonatal screening for congenital adrenal hyperplasia [letter]. Lancet 1982
 May 8;1(8280):1069
- Caielli R; Biavaschi G. [Immunoglobulins (G, A, M) in umbilical cord blood: determination using laser nephelometry]. Quad Sclavo Diagn 1984 Jun;20(2):163-70 [Ita]
- Carson DJ; Roberts G; Campbell SL; Carson NA. Screening for congenital hypothyroidism in Northern Ireland. Ir J Med Sci 1982 Sep;151(9):267-71
- Carta Sorcini M; Fiore L; Tomarchio S; Carta S; Romagnoli C; Curro V; Segni G. Pilot study for congenital hypothyroidism, preliminary report. J Endocrinol Invest 1980 Apr-Jun;3(2):181-4
- Carta Sorcini M; Moschini L; Fiore L; Tomarchio S; Di Irio MG; Gilardi E; Romagnoli C; Curro V; Carta S. Familial thyroxine-binding globulin deficiency detected in a pilot screening program for congenital hypothyroidism. J Endocrinol Invest 1982 Jan-Feb;5(1):21-5
- Carta Sorcini M; Tomarchio S; Fiore L; et al. [Transient hypothyroidism with goiter detected by a neonatal screening program]. Pediatr Med Chir 1980;2(6):791-4 [Ita]
- Cassio A; Bernardi F; Piazzi S; Capelli M; Frejaville E; Villa MP;
 Martelli E; Balsamo A; Salardi S; Merighi R; et al. Neonatal
 screening for cystic fibrosis by dried blood spot trypsin assay.
 Results in 47 127 newborn infants from a homogeneous population. Acta
 Paediatr Scand 1984 Jul;73(4):554-8
- Cassio A; Salardi S; Balsamo A; Piazzi S; Bernardi F; Merighi R;
 Turba E; Cacciari E. [Neonatal screening for congenital
 hypothyroidism. Experience at a Regional Center]. Pediatr Med Chir
 1983 Nov-Dec;5(6):457-63 [Ita]
- Cassio A; Turba E; Balsamo A; Ballestrazzi A; Pozzato R; Marinelli M. In vivo isotope study of the thyroid with 99mTc04- in neonatal congenital hypothyroidism. Eur J Nucl Med 1984;9(8):351-3
- Castro O; Winter WP; Lee TC; Headings VE. Prevalence of alpha-chain variants at birth. Am J Clin Pathol 1981 Jan;75(1):56-9
- Catford JC; Bennet GC; Wilkinson JA. Congenital hip dislocation: an increasing and still uncontrolled disability? Br Med J 1982; 285(6354):1527-30
- Cattaneo G; Ghiotti P; Marra A; et al. [Retinal haemorrhagies: Occasional finding during a neonatal ocular screening]. Minerva Oftalmol 1982;24(1):15-20 [Ita]

- [Causes of congenital hypothyroidism. The Thyroid Study Group]. Tijdschr Kindergeneeskd 1984 Dec;52(6):249-52 [Dut]
- Centa A; Sciarratta G; Spandonari M; et al. [Genetic screening of the newborn. Principles, programs and research]. Pediatr Med Chir 1980; 2(4):455-8 [Ita]
- Chamberlain JM. Which prescriptive screening programmes are worth while? J Epidemiol Community Health 1984 Dec; 38(4):270-7
- Characteristics of infantile hypothyroidism discovered on neonatal screening. J Pediatr 1984 Apr; 104(4):539-44
- Check WA. Genetic counseling--still mostly genetics [news]. JAMA 1980 Jul 25;244(4):315-7
- Chen Rui guan; Qian Da long; Guo Di. Benefit of neonatal screening for phenylketouria. A report of the first case treated in China. Chin Med J 1986;99(6):513-14
- Chen Rui guan; Sun Min; Ni Yuan yu; et al. Neonatal hypothyroidism, phenylketonuria and galactosemia screening in metropolitan Shanghai. Chin Med J 1984;97(1):61-5
- Cleghorn G; Benjamin L; Corey M; et al. Age-related alterations in immunoreactive pancreatic lipase and cationic trypsinogen in young children with cystic fibrosis. J Pediatr 1985;107(3):377-81
- Clemens P. [Screening of newborns: Aspects of differential diagnosis].

 Bundesgesundheitsblatt 1983;26(2):43-5 [Ger]
- Clemens P; Plettner C; Gruttner R. [False positive and false negative results in neonatal screening for hypothyroidism]. Dtsch Med Wochenschr 1983 Mar 25;108(12):467-9 [Ger]
- Clemens P; Plettner C; Gruttner R. [Phenylketonuria screening: False-positive and false-negative results]. Dtsch Med Wochenschr 1983;108(16):627-8 [Ger]
- Clemens P; Plettner C; Gruttner R. [Screening for phenylketonuria without false readings from antibacterial agents]. Off Gesundheitswes 1982 Jun; 44(6):435 [Ger]
- Clemens P; Voltmer C; Plettner C. Interference by antibiotics with neonatal screening for galactosemia. J Pediatr 1986 Oct;109(4):713-4
- Cohen T. Problems in genetic counseling. Prog Clin Biol Res 1982;103 Pt B: 489-96
- Colaco MP; Desai MP; Ajgaonkar AR; Mahadik CV; Vas FE; Bandivdekar AN; Rege CM; Shirodkar VV; Kasi V; Sheth AR. Neonatal screening for hypothyroidism. Indian Pediatr 1984 Sep;21(9):695-700

- Cole DB; Holmes GE. Risk birth registry: A newborn screening model. J Kans Med Soc 1980 Feb;81(2):79-82
- [Conditions for neonatal screening]. Munch Med Wochenschr 1981;123(11):407-9 [Ger]
- Cossu G; Manca M; Pirastru MG; Bullitta R; Bosisio AB; Gianazza E; Righetti PG. Neonatal screening of beta-thalassemias by thin layer isoelectric focusing. Am J Hematol 1982 Sep;13(2):149-57
- Coulombe JT; Shih VE; Levy HL. Massachusetts metabolic disorders screening program. II. Methylmalonic aciduria. Pediatrics 1981; 67(1):26-31
- Cristol P; Des Georges M; Levy A; Sahuc P. [Value of neonatal screening for cystic fibrosis. Evaluation of a neonatal screening program including 34,522 neonates (author's transl)]. Sem Hop Paris 1982 Feb 25;58(8):499-55 [Fre]
- Croner S; Kjellman NIM; Eriksson B; Roth A. IgE screening in 1701 newborn infants and the development of atopic disease during infancy. Arch Dis Child 1982;57(5):364-8
- Crossley JR; Smith PA; Edgar BW; Gluckman PD; Elliott RB. Neonatal screening for cystic fibrosis, using immunoreactive trypsin assay in dried blood spots. Clin Chim Acta 1981 Jun 18;113(2):111-21
- Czernichow P; Vandalem JL; Hennen G. Transient neonatal hyperthyrotropinemia: a factitious syndrome due to the presence of heterophilic antibodies in the plasma of infants and their mothers. J Clin Endocrinol Metab 1981 Aug;53(2):387-93

-				
L)			
_			 	

- Dagenais DL; Courville L; Dagenais MG. A cost-benefit analysis of the Quebec Network of Genetic Medicine. Soc Sci Med 1985;20(6):601-7
- Dailey PA; Baysinger CL; Levinson G; Shnider SM. Neurobehavioral testing of the newborn infant. Effects of obstetric anesthesia. Clin Perinatol 1982;9(1):191-214
- Dalens B; Bezou MJ; Travade P; Raynaud EJ; Coulet M. [Rapid screening for neonatal infection by employing new techniques for the use of the nitroblue tetrazolium test (NBT) (author's transl)]. J Gynecol Obstet Biol Reprod (Paris) 1981;10(1):39-43 [Fre]
- Dandona P; Littlewood JM; Ramdial L; Evans R. Two tier screen for cystic fibrosis [letter]. Lancet 1981 Feb 14;1(8216):380
- Dankert JE; Te Meerman GJ; Knol K; et al. Neonatal screening for cystic fibrosis. Lancet 1986;1(8476):326-8

- Dankert Roelse JE; Meerman GJ; Cornel MC; Knol K; ten Kate LP. Neonatal screening for cystic fibrosis [letter]. Lancet 1986 Apr 5;1(8484):802-3
- Davidson AGF; Wong LTK; Kirby LT; Applegarth DA. Immunoreactive trypsin in cystic fibrosis. J Pediatr Gastroenterol Nutr 1984;3(Suppl.1):S79-S88
- De Filippis V; Fabris C; Bacolla A; Bertino E; Migliardi M; Trapani G; Chillemi C; Mombro M; Mongardi L; Montrucchio F; Garzena E; Costa A. [Screening for congenital hypothyroidism. Results of screening of 20,000 newborn infants]. Minerva Pediatr 1981 Apr 15; 33(7):289-98 [Ita]
- De Magalhaes J. [Neonatal screening (phenylketonuria and hypothyroidism) in Portugal]. J Genet Hum 1981;29(1):33 [Fre]
- De Meyer R. [Neonatal screening of inborn errors of metabolism in Belgium].

 J Genet Hum 1981;29(1):35-8 [Fre]
- Delamonica C; Collombel C; Cotte J; Addis P. Screening for neonatal Duchenne muscular dystrophy by bioluminescence measurement of creatine kinase in a blood sample spotted on paper. Clin Chem 1983;29(1):161-3
- Delange F; Heidemann P; Bourdoux P; et al. Regional variations of iodine nutrition and thyroid function during the neonatal period in Europe. Biol Neonate, Foetal Neonatal Res 1986;49(6):322-30
- Delange F; Illig R; Rochiccioli P; Brock Jacobsen B. Progress report 1980 on neonatal thyroid screening in Europe. Acta Paediatr Scand 1981; 70(1):1-2
- Demarquez JL. [Value of an Apgar score]. Rev Prat 1986;36(6):272-5 [Fre]
- Denaro A; Biondi S; Patane Tropea C; Biondi Zappala M. [Neonatal audiological screenings]. Otorinolaryngologia 1983;33(2):129-40 [Ita]
- Dennis JM; Sheldon R; Toubas P; McCaffee MA. Identification of hearing loss in the neonatal intensive care unit population. Am J Otol 1984 Jan;5(3):201-5
- Derksen Lubsen G. [The diagnosis of galactosemia: screening or clinical diagnosis?] Ned Tijdschr Geneeskd 1980 Jul 26;124(30):1250-6 [Dut]
- Derksen Lubsen G; de Jonge GA; Schopman W. [Mass screening for congenital hypothyroidism in newborn infants in a test area]. Ned Tijdschr Geneeskd 1980 Feb 23;124(8):260-3 [Dut]
- Dhondt JL; Dorche C; Farriaux JP; Courte C. Neonatal screening for congenital adrenal hyperplasia: a pilot study in France. J Inherited Metab Dis 1986;9 Suppl 1:147-51

- Dhondt JL; Leroux B; Farriaux JP; Largilliere C; Leeming RJ. Dihydrobiopterin biosynthesis deficiency. Eur J Pediatr 1983 Dec;141(2):92-5
- Dijkman JH; Penders TJ; Kramps JA; Sonderkamp HJ; van den Broek WG; ter Haar BG. Epidemiology of alpha 1-antitrypsin deficiency in the Netherlands. Hum Genet 1980;53(3):409-13
- Dluholucky S; Bucek M; Hornova V; Hruskovi:c I; Slivka L; Lukac P. [The present state of screening for congenital hypothyroidism in Slovakia]. Cesk Pediatr 1986 May;41(5):283-5 [Slo]
- Dluholucky S; Hornova V; Bucek M; Lukac P. [Results of 3 years of screening for congenital hypothyroidism in the catchment area of the southern Central Slovakia Region]. Cesk Pediatr 1983 Dec;38(12):712-6 [Slo]
- Dluholucky S; Hornova V; Hudec P; Bucek M; Fukal J; Lukac P. [A case of congenital nephrotic syndrome detected during screening for congenital hypothyroidism]. Cesk Pediatr 1982 Apr;37(4):203-5 [Slo]
- Dluholucky S; Hornova V; Hutyrova V. [How to correctly collect a dry drop of blood for screening]. Cesk Pediatr 1982 Sep;37(9):550-1 [Slo]
- Dockeray S; Cahalane SF; Brody M; Mullins C; Cullen MJ. Screening for congenital hypothyroidism in the Republic of Ireland. Br Med J 1980 Dec 6;281(6254):1519-22
- Dockter G; Steyns M; Biro G; Sitzmann FC. [Immunoreactive trypsin in the scrum of normal children and children with gastrointestinal diseases].

 Monatsschr Kinderheilkd 1984 Aug;132(8):589-93 [Ger]
- Dodge JA; Kirk JM; Westwood A. Neonatal screening for cystic fibrosis. Lancet 1986;1(8481):615
- Dodge JA; Ryley HC. Screening for cystic fibrosis [letter]. Arch Dis Child 1983 Apr;58(4):317-8
- Dodge JA; Ryley HC. Screening for cystic fibrosis. Arch Dis Child 1982 Oct;57(10):774-80
- Dominici R; Monaco F; Morano S; Antonozzi I. Immunoreactive trypsin on dried-blood spots as a possible neonatal test for cystic fibrosis.

 (I. Evaluation of the method and preliminary field trial). Ric Clin Lab 1980 Jul-Sep; 10(3):511-9
- **Donovan C.** The challenge of preventive paediatrics. Practitioner 1983; 227(1376):309-11
- Donovan P. Wrongful birth and wrongful conception: the legal and moral issues. Fam Plann Perspect 1984 Mar-Apr;16(2):64-9

- Duara S; Suter CM; Bessard KK; Gutberlet RL. Neonatal screening with auditory brainstem responses: results of follow-up audiometry and risk factor evaluation. J Pediatr 1986 Feb;108(2):276-81
- Dudman NP; Blades BL; Wilcken DE; Aitken JM. Radial immunodiffusion assay of apolipoprotein B in blood dried on filter paper--a potential screening method for familial type II hypercholesterolaemia. Clin Chim Acta 1985 Jul 15;149(2-3):117-27
- Duncan MH; Watterberg KL. Neonatal sepsis screening re hemolytic disease [letter]. J Pediatr 1984 Aug;105(2):337-8
- Dunger DB; Grant DB. Endocrine disorders: Diabetes mellitus, congenital hypothyroidism, and congenital adrenal hyperplasia. Br Med Bull 1986; 42(2):1-7-90
- Durleux Smith A; Jacobson JT. Comparison of auditory brainstem response and behavioral screening in neonates. J Otolaryngol 1985 Feb;14 Suppl 14:47-53
- **Dussault JH.** [Congenital hypothyroidism: A reappraisal]. Union Med Can 1981;110(7):588-93 [Fre]
- Dussault JH; Letarte J; Guyda H; Laberge C. Lack of influence of thyroid antibodies on thyroid function in the newborn infant and on a mass screening program for congenital hypothyroidism. J Pediatr 1980; 96(31):385-9
- Dussault JH; Morissette J. Automated solid-phase assay for estimation of thyroxine binding capacity in a screening program for congenital hypothyroidism [letter]. Clin Chem 1981 Apr;27(4):643
- Dussault JH; Morissette J. Higher sensitivity of primary thyrotropin in screening for congenital hypothyroidism: a myth? J Clin Endocrinol Metab 1983 Apr;56(4):849-52
- Dussault JH; Morissette J; Letarte J; et al. Thyroxine-binding globulin capacity and concentration evaluated from blood spots on filter-paper in a screening program for neonatal hypothyroidism. Clin Chem 1980; 26(3):463-5

E

- Early infant death in sickle cell disease [editorial]. Lancet 1983 May 21; 1(8334):1141-1142
- Eaton AP; Doody DM; Zipf WB; Ackerman JH; Croft C; Porter L. Neonatal hypothyroid screening in Ohio. Ohio State Med J 1982 May;78(5):360-1, 365, 368
- Edwards RW; Hulse JA; Jackson D; Spracklan A; Clayton BE. Programming and running radioimmunoassay in the h-TSH screening procedure using

- dried blood spots and the NE1600 gamma counter on line to the HP9815S calculator. Ann Clin Biochem 1980 May;17(3):122-9
- Eguileor I; Arena J; Aniel A; Rodriguez JL; Galdeano L; Sanjurjo P. [Problems of neonatal screening for hypothyroidism in the very low birth-weight neonate]. An Esp Pediatr 1985 Jun;22(8):551-5 [Spa]
- Endo F; Matsuda I. Screening method for prolidase deficiency. Hum Genet 1981;56(3):349-51
- Enjaume C; Dutau G; Demeurisse P; et al. [Automation of 14 and TSH radioimmunoassay in the neonatal screening for hypothyroidism in South-West France]. Arch Fr Pediatr 1980;37(1):25-8 [Fre]
- Erbe RW. Issues in newborn genetic screening. Birth Defects 1981;17(1):167-79
- Evans RT; Little AJ; Steel AE; Littlewood JM. Satisfactory screening for cystic fibrosis with the BM meconium procedure. J Clin Pathol 1981 Aug;34(8):911-3
- Evert W; Gruettner R. [Hypothyreosis: Screening of newborns]. Z Geburtshilfe Perinatol 1980;184(3):180-6 [Ger]
- Excler JL; Piens MA; Maisonneuve H; et al. [Toxoplasmosis screening in pregnant women and their newborn]. Lyon Med 1985;253(1-2):33-8 [Fre]

-				
r				
•				
_	 			 _

- Faden R; Chwalow AJ; Holtzman NA; Horn SD. A survey to evaluate parental consent as public policy for neonatal screening. Am J Public Health 1982 Dec;72(12):1347-52
- Faden RR; Chwalow AJ; Holtzman NA; Horn SD. A survey to evaluate parental consent as public policy for neonatal screening. Am J Public Health 1982 Dec;72(12):1347-1352
- Faden RR; Holtzman NA; Chwalow AJ. Parental rights, child welfare, and public health: the case of PKU screening. Am J Public Health 1982 Dec;72(12):1396-1400
- Fall M; Ba M; Sarr M; Senghor G. [Neonatal screening for mucoviscidosis in Senegal using the BM-test]. Dakar Med 1983;28(4):713-24 [Fre]
- Farfel MR; Holtzman NA. Education, consent, and counseling in sickle cell screening programs: report of a survey. Am J Public Health 1984 Apr; 74(4):373-5
- Farriaux JP. [Mass neonatal screening. A reality which is sometimes poorly understood]. Arch Fr Pediatr 1980 Jun;37(6):367-70 [Fre]
- Farriaux JP. [Neonatal detection of hypothyroidism. Usefulness and limitations]. Presse Med 1983;12(24):1519-21 [Frc]

- Farriaux JP. [Systematic neonatal screening. Status and organization.

 Apropos of the French experience]. LARC Med 1982 Dcc;2(11):963-7
 [Fre]
- Farriaux JP; Dhondt JL. [Neonatal screening for phenylketonuria]. J Genet Hum 1981 Mar;29(1):3-11 [Fre]
- Farriaux JP; Dhondt JL. [Neonatal screening of congenital hyperplasia of the adrenal glands (21-hydroxylase deficiency)]. Rev Med Liege 1984 May 15;39(10):460-3 [Fre]
- Farriaux JP; Dhondt JL. Newborn thyroid screening [letter]. Am J Dis Child 1982 Nov;136(11):1030
- Farriaux JP; Dhondt JL; Cartigny B. [Screening for neonatal hypothyroidism by estimation of T4 (author's transl)]. Arch Fr Pediatr 1980 Apr;37(4): 227-31 [Fre]
- Faure C; Schmit P; Salvat D. Cost-benefit evaluation of systematic radiological diagnosis of congenital dislocated hip. Pediatr Radiol 1984;14(6):407-12
- Fernhoff PM; Fitzmaurice N; Milner J; McEwen CT; Dembure PP; Brown AL; Wright L; Acosta PB; Elsas LJ 2d. Coordinated system for comprehensive newborn metabolic screening. South Med J 1982 May; 75(5):529-32
- Filetti S; Catanuso A; Lisi E; Pezzino V; Sangiorgio L; Squatrito S; Vigneri R. Measurement of reverse triidothyronine in dried blood spot: methodology and clinical application as a screening procedure for detection of congenital hypothyroidism. Metabolism 1980 Fcb;29(2):141-5
- Fior R; Tamburini P. [Screening with BOEL-test for early detection of hearing defects in infants]. Laryngol Rhinol Otol 1985;64(5):260-2 [Ger]
- Fiorucci G; Milia U; Fedeli L; et al. [Screening for neonatal hypothyroidism: Personal experience]. Arch Med Interna 1983;35(5):401-2 [Ita]
- Firth M; Gardner MD; Hosking G; Wilkinson E. Interviews with parents of boys suffering from Duchenne muscular dystrophy. Dev Med Child Neurol 1983 Aug;25(4):466-71
- Firth MA; Wilkinson EJ. Screening the newborn for Duchenne muscular dystrophy: parents' views. Br Med J [Clin Res] 1983 Jun 18;286(6382): 1933-4
- Fisher DA. Erratum: neonatal thyroid screening [letter]. J Pediatr 1984 Jan;104(1):163-4

- Fisher DA. Second International Conference on Neonatal Thyroid Screening: progress report. J Pediatr 1983 May;102(5):653-4
- Fok TF; Lau SP; Fung KP. Cord blood G-6-PD activity by quantitative enzyme assay and fluorescent spot test in Chinese neonates. Aust Paediatr J 1985;21(1):23-5
- Forrest DC; Wilchen B; Turner G. Screening for cystic fibrosis by a stool trypsin method. Arch Dis Child 1981 Feb; 56(2):151-3
- Foster K; Forbes M; Hayes R; Serjeant GR. Cord blood screening for sickle hemoglobin: evidence against a female preponderance of hemoglobin S. J Pediatr 1981 Jan;98(1):79-81
- Frankenburg WK. To screen or not to screen: congenital dislocation of the hip [editorial]. Am J Public Health 1981 Dec;71(12):1311-1313
- Frey C. [Klinefelter's syndrome: Is an early screening necessary?]
 Monatsschr Kinderheilkd 1986;134(2):78-83 [Ger]
- Frezal J; Briard ML; Saudubray JM. [After PKU and hypothyroidism, what other screening?] J Genet Hum 1981 Mar;29(1):47-57 [Fre]
- Fries F; Lelong Tissier MC; Rolland M; et al. [Neonatal intestinal obstruction due to congenital hypothyroidism]. Arch Fr Pediatr 1980; 37(6):393-5 [Fre]
- Fritsche F; Berndt S. [Diagnostic value and reliability of the newborn hearing-screening: A 7-years investigation]. Hno Prax 1983;8(2):97-100 [Ger]
- Frye Osier HA; Goldstein R; Hirsch JE; Weber K. Early- and middle-AER components to clicks as response indices for neonatal hearing screening. Ann Otol Rhinol Laryngol 1982;91(31):272-6
- Fujimura Y; Kawamura M; Naruse H. A new method of blood galactose estimation for mass screening of galactosemia. Tohoku J Exp Med 1981 Apr;133(4):371-8
- Fujimura Y; Kawamura M; Naruse H. A new mass screening method of detecting UDP-galactose-4-epimerase deficiency. Tohoku J Exp Med 1980 May;131(1):15-22
- Fukushi M; Arai O; Mizushima Y; Namba M; Sato Y; Hayashi H. [Studies of T4 determination by solid-phase radioimmunoassay for mass screening of congenital hypothyroidism]. Horumon To Rinsho 1982 Aug;30(8):889-92 [Jpn]
- Fukushi M; Arai O; Mizushima Y; Takasugi N; Fujieda K; Matsuura N.
 [Neonatal screening for congenital adrenal hyperplasia due to 21hydroxylase deficiency. 1. Enzyme immunoassay of dried blood 17 alpha-

- hydroxyprogesterone and its application to neonatal screening for congenital adrenal hyperplasia]. Nippon Naibunpi Gakkai Zasshi 1986 Jun 20;62(6):683-96 [Jpn]
- Fukuwatari Y. [Mass-screening program for congenital hypothyroidism in Japan (author's transl)]. Horumon To Rinsho 1981 Mar;29(3):277-82 [Jpn]
- Furbetta M; Angius A; Tuveri T; Rosatelli C; Scalas MT; Ximenes A; Falchi AM; Cao A. Diagnosis of the beta 0 thalassemia trait at birth. Hemoglobin 1981;5(3):217-29
- Furmaier R; Staudt F. [Reconstruction in coronal and sagittal planes of CT scans compared with two-dimensional echo-encephalography in neonates and infants (author's transl)]. ROFO 1982 Jul;137(1):37-42 [Ger]

4	_
	_

- Galacteros F; Kleman K; Caburi Martin J; Beuzard Y; Rosa J; Lubin B. Cord blood screening for hemoglobin abnormalities by thin layer isoelectric focusing. Blood 1980 Dec;56(6):1068-71
- Galanello R; Diana G; Furbetta M; Angius A; Melis MA; Rosatelli C;
 Cao A. alpha-Thalassaemia in Sardinian infants. J Med Genet 1980 Oct;
 17(5):357-62
- Galasko CS; Galley S; Menon TJ. Detection of congenital dislocation of the hip by an early screening program, with particular reference to false negatives. Isr J Med Sci 1980 Apr;16(4):257-9
- Garrick MD; Orfanos AP; Rogers L; Naylor EW; Guthrie R. A simple screening test for reduced glutathione in filter paper spots of blood. J Pediatr 1981 Feb;98(2):265-7
- Gaston MH; Verter JI; Woods G; Pegelow C; Kelleher J; Presbury G; Zarkowsky H; Vichinsky E; Iyer R; Lobel JS; et al. Prophylaxis with oral penicillin in children with sickle cell anemia. A randomized trial. N Engl J Mcd 1986 Jun 19;314(25):1593-9
- Gendrel D; Feinstein MG; Grenier J; et al. Falsely elevated serum thyrotropin (TSH) in newborn infants: Transfer from mothers to infants of a factor interfering in the TSH radioimmunoassay. J Clin Endocrinol Metab 1981;52(1):62-5
- Genetic counseling and prevention of birth defects. JAMA 1982 Jul 9; 248(2):221-224
- Gerald PS. Regionalization of genetic services. Birth Defects 1983; 19(5):127-33
- Giaquinto C; Vanin M; Anglani F; Errico G; Ruga E; D'Elia R. A new screening test for rotavirus infection. Infection 1985 Nov-Dec; 13(6):260-2

- Gibson LE. Screening of newborns for cystic fibrosis [editorial]. Am J Dis Child 1980 Oct;134(10):925-6
- Giguere R; Auray Blais C; Draper P; Lemieux B. Dict and medications giving positive ninhydrin reactions on TLC in a newborn urinary screening program. Clin Biochem 1980 Jun;13(3):103-5
- Gilbert GL. Prevention of vertical transmission of hepatitis B. The place of routine screening in antenatal care, and the case for immunization of infants at risk. Med J Aust 1984 Aug 18;141(4):213-6
- Gitzelmann R. Rationales for genetic screening. Prog Clin Biol Res 1982;103 Pt B:425-36
- Gitzelmann R. Why we should not screen our newborns for cystic fibrosis. Helv Paediatr Acta 1981;36(6):493-4
- Giugliani R; Dutra JC; Pereira ML; Rotta N; Drachler M de L; Ohlweiller L; Pina Neto JM; Pinheiro CE; Breda DJ. GMl gangliosidosis: clinical and laboratory findings in eight families. Hum Genet 1985;70(4):347-54
- Glader BE. Screening for anemia and erythrocyte disorders in children. Pediatrics 1986 Aug;78(2):368-9
- Glorieux J; Dussault JH; Letarte J; Guyda H; Morissette J. Preliminary results on the mental development of hypothyroid infants detected by the Quebec Screening Program. J Pediatr 1983 Jan;102(1):19-22
- Golding J; Halsum M; Carswell F. Reduced morbidity in cystic fibrosis. Lancet 1986;1(8472):104-5
- Gottfried EL. Screening for beta-thalassaemia [letter]. Lancet 1980 Oct 11;2(8198):806
- Gotz M; Stur O; Zach M. [New data on cystic fibrosis (mucoviscidosis)].
 Osterr Arzteztg 1982;37(13-14):904-8 [Ger]
- Grant DB; Hulse JA. Screening for congenital hypothyroidism [editorial].

 Arch Dis Child 1980 Dec;55(12):913-4
- Grant GA; Carson DJ; Reid MM; Hutchinson JM. Congenital hypothyroidism missed on screening. Arch Dis Child 1986 Feb;61(2):189-90
- Grenier J; Canlorbe P; Roger M; Scholler R. [Screening for congenital hypothyroidism in 26,282 newborns]. Pathol Biol (Paris) 1980; 28(6):387-8 [Fre]
- Griffiths KD; Raine DN; Mann JR. Neonatal screening for sickle haemoglobinopathics in Birmingham. Br Mcd J [Clin Res] 1982 Mar 27; 284(6320):933-5

- Griffiths KD; Virdi NK; Rayner PH; Green A. Noonatal screening for congenital hypothyroidism by measurement of plasma thyroxine and thyroid stimulating hormone concentrations. Br Med J [Clin Res] 1985 Jul 13;291(6488):117-20
- Grimm T. [Newborn screening for Duchenne muscular dystrophy (author's transl)]. Monatsschr Kinderheilkd 1981 Jul;129(7):414-7 [Ger]
- Grobe H. [Screening studies in newborn infants]. Monatsschr Kinderheilkd 1983 Nov;131(11):806-9 [Ger]
- Grosse KP; Busanny W; Fleckenstein B. [Results of precise screening for prenatal cytomegalovirus and BK virus infections in newborn and young infants. I. CMV infections]. Monatsschr Kinderheilkd 1980 May; 128(5):267-8 [Ger]
- Grossman LK. On newborn sickle cell screening in NYC [letter]. Am J Public Health 1983 Oct;73(10):1216-7
- Grossman LK; Holtzman NA; Charney E; Schwartz AD. Neonatal screening and genetic counseling for sickle cell trait. Am J Dis Child 1985 Mar; 139(3):241-4
- Grover R; Newman S; Wethers D; Anyane Yeboa K; Pass K. Newborn screening for hemoglobinopathics: the benefit beyond the target. Am J Public Health 1986 Oct;76(10):1236-7
- Grover R; Shahidi S; Fisher B; Goldberg D; Wethers D. Current sickle cell screening program for newborns in New York City, 1979-1980. Am J Public Health 1983 Mar;73(3):249-52
- Gruters AL; Allemand D; Beyer P; et al. [Newborn screening for hypothyroidism in Berlin 1978-1982]. Monatsschr Kinderheilkd 1983; 131(2):100-5 [Ger]
- Gruttner R; Clemens P; Koepp P; Held KH; Plettner C; Stern M.

 [Neonatal screening for mucoviscidosis using the BM-test-meconium].

 Monatsschr Kinderheilkd 1985 Jan;133(1):54-6 [Ger]
- Gustavson KH; Zetterstrom R. [Mctabolic screening of nconates].

 Lakartidningen 1982;79(4):191-3 [Swc]
- Guthrie R; Susi A. Laboratory confirmation of phenylketonuria. Combined paper chromatography and "inhibition assay." Padiatr Padol 1982;17(2): 165-9

H

Hagenfeldt L; Lundberg K; Larsson A; Thuresson M. [Information from the PKU-section - report of activities 1981]. Jordemodern 1982 Sep; 95(9):287-90 [Swc]

- Hahn HB Jr; Brown LO; Hillis A; Spiekerman AM; Levine G. Broast feeding and neonatal screening for congenital hypothyroidism. Tex Med 1986 Jan;82(1):46-7
- Haibach H; Woodruff CW. Neonatal screening for metabolic diseases in Missouri. Mo Med 1982 Sep;79(9):615-20
- Haibach H; Woodruff CW; Harris DJ; Baumgartner JH. Galactosemia: Progress in neonatal screening in Missouri. Mo Med 1986;83(5):271-4
- Halberg F; Cornelissen G; Bingham C; Tarquini B; Mainardi G; Cagnoni M; Panero C; Scarpelli P; Romano S; Marz W; et al. Neonatal monitoring to assess risk for hypertension. Postgrad Med 1986 Jan; 79(1):44-6
- Hanna CE; Krainz PL; Skeels MR; Miyahira RS; Sesser DE; LaFranchi SH.

 Detection of congenital hypopituitary hypothyroidism: ten-year
 experience in the Northwest Regional Screening Program. J Pediatr 1986
 Dec;109(6):959-64
- Hanshaw JB. On deafness, cytomegalovirus, and neonatal screening. Am J Dis Child 1982 Oct;136(10):886-7
- Hansteen IL; Varslot K; Steen Johnsen J; Lang-ard S. Cytogenetic screening of a new-born population. Clin Genet 1982 May;21(5):309-14
- Harada T; Nose O; Maki I; Kai H; Yabuuchi H. [The problems of treatment of neonatal cretinism detected by mass screening (author's transl)]. Horumon To Rinsho 1981 Mar;29(3):327-8 [Jpn]
- Harper JI; Smith P; Dickey D; Broussard ER. Screening and assessment of psychosocial function in a private pediatric service. Infant Ment Health J 1982 Fall;3(3):199-208
- Harris P; Dreyfus NG. Newborn thyroid screening in a municipal hospital.

 Am J Dis Child 1982 Mar;136(3):248-50
- Hata N; Ito M; Mizuta H; Nose O; Miyai K. Enzymc immunoassay of thyroxin-binding globulin in dried blood samples on filter paper. Clin Chem 1983 Jul;29(7):1437-40
- Hata N; Miyai K; Ito M; Endo Y; Iijimi Y; Mizuta H; Amino N; Nose O; Harada T. Enzyme immunoassay of free thyroxin in dried blood samples on filter paper. Clin Chem 1985 May;31(5):750-3
- Hayashi A; Wada Y. Neonatal mass-screening of hemoglobin variants using dried blood on filter paper. Jpn J Hum Genet 1985;30(2):103-4
- Hayashi M; Ezaki Y; Ikumi S; Noda M; Yamashita F. [Newborn screening for hypothyroidism--the evaluation of methods (author's transl)].

 Horumon To Rinsho 1981 Mar;29(3):285-8 [Jpn]

- Headings V. Counselling in a hospital-based newborn screening service.

 Patient Couns Health Educ 1980 2d Quart;2(2):80-3
- Headings VE; Winter W; Reindorf C; Castro O. Sickle Lepore hemoglobin identified in a black American infant. Am J Pediatr Hematol Oncol 1983 Fall;5(3):259-64
- Healy A. Screening for disabilities [editorial]. West J Med 1985 Sep; 143(3):379-80
- Heard GS; Secor McVoy JR; Wolf B. A screening method for biotinidase deficiency in newborns. Clin Chem 1984 Jan;30(1):125-7
- Heard GS; Wolf B; Jefferson LG; Weissbecker KA; Nance WE; McVoy JR; Napolitano A; Mitchell PL; Lambert FW; Linyear AS. Nconatal screening for biotinidase deficiency: results of a 1-year pilot study. J Pediatr 1986 Jan;108(1):40-6
- Hearn TL; Hannon WH. Interlaboratory surveys of the quantitation of thyroxin and thyrotropin (thyroid-stimulating hormone) in dried blood spot specimens. Clin Chem 1982;28(10):2022-5
- Hecht F. Screening people of Jewish origin for Tay-Sachs disease carriers.

 Ariz Med 1981 Jan;38(1):21-3
- Heeley AF. Screening for cystic fibrosis [letter]. J Pediatr 1983 Nov; 103(5):836
- Heeley AF; Heeley ME; King DN; Kuzemko JA; Walsh MP. Screening for cystic fibrosis by died blood spot trypsin assay. Arch Dis Child 1982 Jan;57(1):18-21
- Hellsing K; Barrljung K; Ceder O; Kollberg H. Mcconium screening for cystic fibrosis. An eight-year follow-up study. Acta Paediatr Scand 1982 Sep;71(5):827-32
- Henssge EJ. [Methods and significance of screening examination of the hip joint in infants, toddlers and preschool children]. Off Gesundheitswes 1982;44(3):161-4 [Gcr]
- Henthorn J; Anionwu E; Brozovic M. Screening cord blood for sickle haemoglobinopathies in Brent. Br Med J [Clin Res] 1984 Aug 25; 289(6443):479-80
- Herrera AJ; Hsiang YH. Comparison of various methods of blood sugar screening in newborn infants. J Pediatr 1983 May;102(5):769-72
- Hesse V; Streckenbach B; Weber A; Strube M. [Screening for congenital hypothyroidism in Europe]. Radiol Diagn 1982;23(1):92-3 [Gcr]
- Higurashi M. [Screening for chromosome abnormalities, especially for cases with sex chromosomal mosaics]. No To Hattatsu 1983;15(2):127-32 [Jpn]

- Hikita Y. [Reevaluation of recalled infants by neonatal mass screening for congenital adrenal hyperplasia due to 21-hydroxylase deficiency. Diagnostic value of pregnanetriolone in a single urine specimen using glass capillary gas chromatography]. Nippon Naibunpi Gakkai Zasshi 1985 Mar 20;61(3):197-219 [Jpn]
- Hitzeroth HW; t'Hof JO. Screening for hypothyroidism in south africa [letter]. S Afr Med J 1981 Dec 5;60(23):880-1
- Hjalmarson O. Epidemiology and classification of acute, neonatal respiratory disorders. A prospective study. Acta Paediatr Scand 1981; 70(6):773-83
- Hnikova O; Dienstbier Z. [Results of screening for congenital hypothyroidism in trials conducted by several facilities in Czechoslovakia]. Cesk Pediatr 1983 May;38(5):268-70 [Cze]
- Hnikova O; Kracmar P. [Inborn defect of thyroxine binding globulin (TBG) diagnosed during screening for congenital hypothyroidism]. Cesk Pediatr 1983 Jul;38(7):416-8 [Czc]
- Hnikova O; Kracmar P. The thyroxine screening index in congenital hypothyroidism screening. Acta Paediatr Hung 1985;26(1):49-52
- Hnikova O; Kracmar P; Ticha H; Wiererova O. [Early detection of congenital hypothyroidism in paediatric practice]. Cesk Pediatr 1980; 35(7):345-7 [Cze]
- Hoerst M; Hohlweg Majert P; Kachel W; et al. [Screening for hypothyroidism in newborns: Results and validity of FTsub 4-RIA]. Nuc Compact 1982;13(1):16-20 [Ger]
- Hoffman GL; Laessig RH; Hassemer DJ; Makowski ER. Dual-channel continuous-flow system for determination of phenylalanine and galactose: application to newborn screening. Clin Chem 1984 Feb; 30(2):287-90
- Hofman H; Czochanska J. [Evaluation of the neurological status of newborn infants by screening methods]. Pediatr Pol 1980 Aug;55(8):927-36 [Pol]
- Holmes GE; Hassanein RS. The KIDS chart. A simple, reliable infant development screening tool. Am J Dis Child 1982 Nov;136(11):997-1001
- Holmes LB; Cann C; Cook C. Examination of infants for both minor and major malformations to evaluate for possible teratogenic exposures. Prog Clin Biol Res 1985;163B:59-63
- Holtzman C; Slazyk WE; Cordero JF; Hannon WH. Descriptive epidemiology of missed cases of phenylketonuria and congenital hypothyroidism. Pediatrics 1986 Oct;78(4):553-8
- Holtzman NA. The impact of the federal cutback on genetic services. Am J Med Genet 1983 Jun;15(2):353-71

- Holtzman NA. Pitfalls of newborn screening (with special attention to hypothyroidism): when will we ever learn? Birth Defects 1983; 19(5):111-20
- Holtzman NA. Routine screening of newborns for cystic fibrosis: not yet. Pediatrics 1984 Jan;73(1):98-9
- Holtzman NA; Faden R; Chwalow AJ; Horn SD. Effect of informed parental consent on mothers' knowledge of newborn screening. Pediatrics 1983 Dec;72(6):807-12
- Holtzman NA; Leonard CO; Farfel MR. Issues in antenatal and neonatal screening and surveillance for hereditary and congenital disorders.

 Annu Rev Public Health 1981;2:219-51
- Holtzman NA; McCabe ER; Cunningham GC; Berry HK. Screening for phenylketonuria [letter]. N Engl J Med 1981 May 21;304(21):1300-1
- Honour J. Biochemical aspects of congenital adrenal hyperplasia. J Inherited Metab Dis 1986;9 Suppl 1:124-34
- Hopfner B. [Mcthodological study of different qualities of filter paper in the radioimmunological determination of thyrotropin in newborn infants]. J Clin Chem Clin Biochem 1982 Dec;20(12):915-20 [Ger]
- Horn ME; Dick MC; Frost B; Davis LR; Bellingham AJ; Stroud CE; Studd JW. Neonatal screening for sickle cell diseases in Camberwell: results and recommendations of a two year pilot study. Br Med J [Clin Res] 1986 Mar 15;292(6522):737-40
- Hornchen H; Betz R; Kotlarek F; Roebruck P. Microprocessor-based long term cardiorespirography. II. Status evaluation in term and premature newborns. J Perinat Med 1983;11(1):32-42
- Hubbard R; Henifin MS. Genetic screening of prospective parents and of workers: some scientific and social issues. Int J Health Serv 1985; 15(2):231-51
- Hubl W; Feher T; Rohde W; Dorner G; Taubert H; Freymann E. Enzyme immunoassay of 17-hydroxyprogesterone in plasma, microfilter paper blood and saliva of newborns, children and patients with congenital adrenal hyperplasia. Endokrinologie 1982 Jun;79(2):165-72
- Huisjes HJ; Lok Meijer TY; Touwen BCL; Olinga AA. Neurological sequelac in high- and low-optimality obstetrics. Gynecol Obstet Invest 1983; 15(5):318-22
- Huisman THJ; Cameron BF; Castro O; Mentzer WC. Recommendations for uniform procedures for neonatal screening. Hemoglobin 1985;9(5):543-4
- Hulse JA; Grant DB; Clayton BE; Lilly P; Jackson D; Spracklan A; Edwards RW; Nurse D. Population screening for congenital hypothyroidism. Br Mcd J 1980 Mar 8;280(6215):675-8

- Hulse JA; Grant DB; Jackson D; Clayton BE. Growth, development, and reassessment of hypothyroid infants diagnosed by screening. Br Med J 1982;284(6327):1435-7
- Hummer L; Munkner T; Sorensen SS; et al. [Screening for congenital hypothyroidism in Denmark. I. Determination of TSH levels in all neonates for a two-year period]. Ugeskr Laeg 1982;144(14):1001-4 [Dan]
- Hummer L; Munkner T; Srensen SS; Brandt NJ; Jacobsen BB. Nation-wide TSH-screening with low recall rate. Laboratory results of a two-year study. Scand J Clin Lab Invest 1982 Feb;42(1):49-55
- Hyanek J. Chromatographic screening for cystinuria among newborns. J Inherited Metab Dis 1980;3(3):77-8
- Hyde ML; Riko K; Corbin H; Moroso M; Alberti PW. A neonatal hearing screening research program using brainstem electric response audiometry. J Otolaryngol 1984 Feb; 13(1):49-54

- Il'in BN. [Physical development indices of children and adolescents in genetic monitoring]. Genetika 1983 Jun;19(6):1024-32 [Rus]
- Iowa neonatal screening: a progress report. J Iowa Mcd Soc 1981 Dcc; 71(12):540-1
- Irie M. [Cretinism--its mass-screening (author's transl)]. Horumon To Rinsho 1981 Mar;29(3):283-4 [Jpn]

_	
•	

- Jacobs S; Peterson L; Thompson L; Tukey D; Paine, Saunders S; Hedlund B; Smith C 2d. Newborn screening for hemoglobin abnormalities. A comparison of methods. Am J Clin Pathol 1986 Jun;85(6):713-5
- Jacobson JT; Mencher GT. Intensive care nursery noise and its influence on newborn hearing screening. Int J Pediatr Otorhinolaryngol 1981 Jan; 3(1):45-54
- Jacobson JT; Morehouse CR. A comparison of auditory brain stem response and behavioral screening in high risk and normal newborn infants. Ear Hear 1984 Jul-Aug;5(4):247-53
- Jahrig D; Julich WD; Heinrich U; Pietsch P; Kombke B. [Hepatitis B screening in pregnant patients and their children--a Greifswald perinatal-pediatric study]. Zentralbl Gynakol 1986;108(12):730-8 [Ger]

- Jenkins MB. Newborn metabolic screening in Minnesota. II. PKU and galactosemia. Minn Med 1981 Jul;64(7):429-32, 398
- Jenkins MB; Foley MH. Newborn metabolic screening in Minnesota--I. Congenital hypothyroidism. Minn Med 1980 Sep;63(9):645-8
- Jinks DC; Guthrie R; Naylor EW. Simplified procedure for producing Bacillus subtilis spores for the Guthrie phenylketonuria and other microbiological screening tests. J Clin Microbiol 1985 May;21(5):826-9
- John R; Woodhead JS. An automated immunoradiometric assay of thyrotrophin (TSH) in dried blood filter paper spots. Clin Chem Acta 1982 Nov 10; 125(3):329-40
- Johnsen NJ; Bagi P; Elberling C. Evoked acoustic emissions from the human ear. III. Findings in neonates. Scand Audiol 1983;12(1):17-24
- Johnson CA; Snyder MS; Weaver RL. Effects of fresh frozen plasma infusions on coagulation screening tests in neonates. Arch Dis Child 1982 Dec;57(12):950-2
- Johnson DF; Woodhams L; Katzenmeyer M. Newborn screening and intervention in the context of health care. Infant Ment Health J 1982 Summer; 3(2):97-105
- Johnson ML; Rumack CM; Mannes EJ; Appareti KE. Detection of neonatal intracranial hemorrhage utilizing real-time and static ultrasound. JCU 1981 Oct;9(8):427-33
- Jones JD. Screening tests for inherited metabolic diseases. Mayo Clin Proc 1984 May;59(5):347-9
- Jorda A; Portoles M. Priority of screening for neonatal hypothyroidism in Spain. Eur J Pediatr 1982;138(4):359
- Juillard E. [Neonatal screening in Switzerland]. J Genet Hum 1981;29(1):31-2 [Fre]

к_____

- Kalra A; Jain AK; Hazra DK; Shukla AK; Mathur DD; Dayal RS. Thyroid screening in newborns. Indian Pediatr 1985 Feb;22(2):131-6
- Karathanos A; Giacoia GP. Should Oklahoma screen newborns for galactosemia? J Okla State Med Assoc 1981 Jun;74(6):169-72
- Karolyi G; Andreka B. First results of a regional neonatal thyroid screening programme in Hungary [letter]. Eur J Pediatr 1984 Nov; 143(1):76
- Kelly S. Galactosemia identified in newborn screening program; clinical and biochemical characteristics. NY State J Med 1980 Nov;80(12):1836-9

- Kelly, Thaddeus E. Genetic screening--eugenics--ethics. In: <his> Clinical Genetics and Genetic Counseling. Chicago: Year Book Medical Publishers, 1980. pp. 262-277
- Kenny D; Ward OC; Mulhern B; Rashied AA. Failure of cord serum cholesterol and betalipoprotein as screening tests for familial hyperlipoproteinaemia. Ir J Med Sci 1984 Mar; 153(3):100-5
- Kerisit J; Senecal J; Fisselier MP; Cormier M; Defawe G; de Queiroz F. [The extemporaneous histological examination of the umbilical cord. A reliable procedure for evaluating the risk of neo-natal infection (author's transl)]. J Gynecol Obstet Biol Reprod (Paris) 1981; 10(1):45-9 [Fre]
- Kessler S; Kessler H; Ward P. Psychological aspects of genetic counseling. III. Management of guilt and shame. Am J Mcd Genet 1984 Mar;17(3):673-97
- Khalifa M; Prehu MO; Galacteros F; Meziane A; Riou J; Godard C; Setaoui A; Rosa R; Rosa J. [Screening for hemoglobinopathies and G6PD deficiencies in Morocco]. Nouv Rev Fr Hematol 1986;28(4):245-7 [Fre]
- Kim HC. Laboratory identification of inherited hemoglobinopathies in children. Clin Pediatr (Phila) 1981 Mar;20(3):161-71
- Kimpen J; Bosmans E; Janssen J; et al. Screening for Ureaplasma urealyticum infections in the neonate and the association with prematurity. Eur J Obstet Gynecol Reprod Biol 1986;22(1-2):53-60
- Kind C. [What does neutropenia tell about the existance of sepsis in the newborn? Predictive value of clinical laboratory tests]. Helv Paediatr Acta 1986 Oct;41(4):277-89 [Ger]
- Kirby LT; Norman MG; Applegarth DA; Hardwick DF. Screening of newborn infants for galactosemia in British Columbia. Can Med Assoc J 1985 May 1;132(9):1033-5
- Kirkman HN; Carroll CL; Moore EG; Matheson MS. Fifteen-year experience with screening for phenylketonuria with an automated fluorometric method. Am J Hum Genet 1982;34(5):743-52
- Kitagawa T. Neonatal mass-screening for inborn errors of metabolism in Japan. Acta Paediatr Jpn 1982;24(1):83-90
- Kitagawa T; Owada M; Sakiyama T; et al. Experience and problems of newborn mass screening for inborn errors of metabolism in Japan. Acta Paediatr Jpn 1981;23(1):24-34
- Kjellman NI; Tilling B. [IgE screening of newborn infants in theory and practice]. Nord Med 1986;101(5):150-1, 165 [Swe]

- Klabuschnigg A; Steinert I; Weissenbacher G. [Neurological development screening of high risk infants in an Austrian provincial district (author's transl)]. Wien Med Wochenschr 1981 Sep 30;131(18):449-52 [Ger]
- Klein RZ. Infantile hypothyroidism then and now: the results of neonatal screening. Curr Probl Pediatr 1985 Jan;15(1):1-58
- Klett M. [Directions for TSH screening of newborn infants]. Dtsch Med Wochenschr 1985 Sep 13;110(37):1423-30 [Ger]
- Klett M; Muller M; Schonberg D; Gruttner R; Clemens P. [False-positive and false-negative results in the screening of newborns for hypothyroidism (letter)]. Dtsch Med Wochenschr 1983 Dec 23;108(51-52):1984-5 [Ger]
- Klett M; Schonberg D. [Neonatal screening for hypothyroidism in the Federal Republic of Germany (author's transl)]. Dtsch Med Wochenschr 1981 Jan 2;106(1):6-12 [Ger]
- Klisic P; Rakic D; Pajic D. Triple prevention of congenital dislocation of the hip. J Pediatr Orthop 1984;4(6):759-61
- Knight PJ; Clatworthy RW Jr. Screening for latent malformations: cost effectiveness in neonates with correctable anomalies. J Pediatr Surg 1982 Apr;17(2):123-9
- Knoppers BM. Genetic information and the law: constraints, liability and rights. Can Med Assoc J 1986 Dec 1;135(11):1257-9
- Koch R; Friedman EG. Accuracy of newborn screening programs for phenylketonuria. J Pediatr 1981 Feb;98(2):267-9
- Koffigan M; Fruchart JC; Dhondt JL; Moschetto Y; Farriaux JP.

 Apolipoprotein B measurement in blood spotted on paper (a screening test for familial type II A hyperlipoproteinemia. Ric Clin Lab 1982 Jan-Mar; 12(1):169-72
- Kollberg H. Diagnosis and main clinical features of cystic fibrosis with special attention to screening procedures. Acta Paediatr Scand 1982; 71(Suppl.301):15-25
- Kovar I; Mayne P; Barltrop D. Plasma alkaline phosphatase activity: A screening test for rickets in preterm neonates. Lancet 1982; 1(8267):308-10
- Krauss JS; Fadel HE; Squires J; Baisden CR. Detection of fetal-maternal hemorrhage in a mother with sickle trait and hereditary persistence of fetal hemoglobin. Transfusion 1983 Nov-Dec;23(6):530-1
- Kreimberg CE. [Study on prevention of phenylketonuria and neonatal hypothyroidism in newborn children]. Neuropsiquiatria Salud Ment 1982 Sep-Dec;13(3):21-3 [Spa]

- Krieger I; Nigro M; Taqi Q. Screening for metabolic disease in a metropolitan hospital. Am J Dis Child 1982 Feb;136(2):125-8
- Krieger IE; Nigro M; Sarnaik A; Taqi Q. Screening of high risk infants for metabolic disease in a metropolitan hospital. J Inherit Metab Dis 1981;4(2):81-2
- Kundrat J; Ranochova S. [Early diagnosis of congenital hip dysplasia in Ostrava 1977-1983]. Acta Chir Orthop Traumatol Cech 1985 Jul;52(4): :332-5 [Cze]
- Kuroda Y; Watanabe T; Ito M; et al. Genetic heterogeneity of histidinemia detected by screening newborn infants in Japan. Jpn J Hum Genet 1985;30(4):287-95
- Kurz C. [Better knowledge of genetic counseling]. Soins 1983 Apr;(403):2 [Fre]
- Kusuda S; Soma T; Ozasa Y; Hase Y; Tsuruhara T. [Pilot study on mass screening for cretinism by TSH-EIA (sandwich method)]. Horumon To Rinsho 1983 Oct;31(10):1011-4 [Jpn]
- Kuyper P. Audiometry and the new-born. Audiology 1981;20(6):530-3
- Kuzemko JA. Screening, early neonatal diagnosis and prenatal diagnosis. J R Soc Med 1986;79 Suppl 12:2-5

L_____

- LaFranchi S. Newborn screening for congenital hypothyroidism. Pediatr Ann 1980 Oct;9(10):390-8
- LaFrancli SH; Hanna CE; Krainz PL; Skeels MR; Miyahira RS; Sesser DE. Screening for congenital hypothyroidism with specimen collection at two time periods: results of the Northwest Regional Screening Program. Pediatrics 1985 Nov;76(5):734-40
- Lambotte C. [Current status of neonatal screening on the national and regional level]. Rev Med Liege 1984 May 15;39(10):413-5 [Fre]
- Lambotte C. [Noonatal screening and its criteria]. Rev Med Liege 1984 May 15;39(10):410-2 [Fre]
- Lambotte C; Schoos Barbette S; Dodinval Versie J. [Screening for mucoviscidosis by meconium analysis. 7 years' experience]. J Genet Hum 1981 Mar;29(1):85-92 [Frc]
- Lambotte C; Schoos Barbette S; Dodinval Versie J. [Screening for mucoviscidosis using meconium analysis. 7 years' results (1973-1980)]. Rev Med Liege 1984 May 15;39(10):448-50 [Fre]

1:3

- Laricheva IP; Chulkova VI; Sirnova TA; Lenni VG. [Radioimmunoassay for screening studies in the newborn]. Mcd Radiol (Mosk) 1983 Aug; 28(8):31-4 [Rus]
- Larsson A; Hagenfeldt L; Westphal O; et al. [Screening for congenital hypothyroidism. Report of a national experimental research study]. Lakartidningen 1982;79(4):221-4 [Swe]
- Larsson A; Ljunggren JG; Ekman K; Nilsson A; Olin P. Screening for congenital hypothyroidism. I. Laboratory results of a pilot study based on dried blood samples collected for PKU screening. Acta Paediatr Scand 1981 Mar;70(2):141-6
- Larsson A; Ljunggren JG; Ekman K; Nilsson A; Olin P; Bodeg~ard G. Screening for congenital hypothyroidism. II. Clinical findings in infants with positive screening tests. Acta Paediatr Scand 1981 Mar; 70(2):147-53
- Larsson A; Ljunggren JG; Lundberg K. TSH and thyroxine in stored neonatal filter-paper blood samples from patients with congenital hypothyroidism. Acta Paediatr Scand 1982;71(1):39-41
- Law AS; Craven EM; Sarafidis EH. Screening of newborns for hemoglobinopathies: results in 5,484 patients. Del Med J 1985 Mar; 57(3):161-4
- Lazarus JH; John R; Ginsberg J; et al. Transient neonatal hyperthyrotrophinaemia: A serum abnormality due to transplacentally acquired antibody to thyroid stimulating hormone. Br Med J 1983; 286(6365):592-4
- Lebovitz RM; Pauli RM; Laxova R. Delayed diagnosis in congenital adrenal hyperplasia. Need for newborn screening. Am J Dis Child 1984 Jun; 138(6):571-3
- Leck I. An epidemiological assessment of neonatal screening for dislocation of the hip. J R Coll Physicians Lond 1986 Jan;20(1):56-62
- Lehmann ECH; Street DG. Neonatal screening in Vancouver for congenital dislocation of the hip. Can Med Assoc J 1981;124(8):1003-8
- Leland D; French ML; Kleiman MB; Schreiner RL. The use of TORCH titers.
 Pediatrics 1983 Jul;72(1):41-3
- Lemieux B; Auray Blais C; Giguere R; Shapcott, D. Prolidase deficiency: detection of cases by a newborn urinary screening programme. J Inherited Metab Dis 1984;7 Suppl 2:145-6
- Levi H. Early detection of hearing loss in infants by auditory nerve and brain stem responses. Audiology 1983 Mar-Apr;22(2):181-8
- Levine DH; Meyer HB. Newborn screening for ABO hemolytic disease. Clin Pediatr (Phila) 1985 Jul;24(7):391-4

- Levine GD; Therrell BL Jr. Second testing for hypothyroidism [letter]. Pediatrics 1986 Aug;78(2):375-6
- Levy HL; Mitchell ML. The current status of newborn screening. Hosp Pract [Off] 1982 Jul;17(7):89-97
- Levy HL; Mitchell ML; Ridley SE. Newborn screening [letter]. Pediatrics 1984 Mar;73(3):417
- Levy HL; Simmons JR; MacCready RA. Stability of amino acids and galactose in the newborn screening filter paper blood specimen. J Pediatr 1985 Nov;107(5):757-60
- Li AM; Lee FT; Todd D. The screening of Chinese cord blood for haemoglobinopathies. Hum Hered 1982;32(1):62-70
- Libert J. Diagnosis of lysosomal storage diseases by the ultrastructural study of conjunctival biopsies. Pathol Annu 1980;15(Pt 1):37-66
- Lie SO; Waarum AS; Waaler AL; Olafsen MK; Kase F; Skjelkvale L; Edminson P. [A nation-wide screening program for congenital hypothyroidism in Norway]. Tidsskr Nor Lacgeforen 1985 Jan 10;105(1): 15-8 [Nor]
- Lindback T; Lie SO. [Screening of congenital hypothyroidism. An analysis of children diagnosed in 1965-74]. Tidsskr Nor Lacgeforen 1985 Jan 10; 105(1):28-30 [Nor]
- Lippman-Hand A. Communication and decision making in genetic counseling.

 Prog Clin Biol Res 1982;103 Pt B:511-9
- Lison M; Sack J; Katznelson D. Early detection of congenital nephrotic syndrome of finnish type by the neonatal hypothyroid screening program [letter]. Int J Pediatr Nephrol 1983 Mar;4(1):61
- Liu Shen ru; Zuo Qi hua. Newborn screening for phenylketonuria in eleven districts. Chin Med J 1986;99(2):113-8
- Lobel JS; Cameron BF; Johnson E; Smith D. The value of screening umbilical cord blood for hemoglobinopathy. Ohio State Med J 1984 Feb; 80(2):140-2
- Loke HL; Lun KC. Reflotest hypoglycemie--a more sensitive screening method for neonatal hypoglycaemia. Singapore Med J 1986 Aug;27(4):321-5
- Lombardo G; De Luca F; Gemelli M; Trimarchi F. [Clinical findings during a 3-year experience in screening newborn infants for congenital hypothyroidism]. Minerva Pediatr 1981 Apr 30;33(8):353-6 [Ita]
- Looye A; van der Vis Melsen MJ; Schreuder CH. Value of the determination of "true' free thyroxine in screening the newborn with possible congenital TBG-deficiency for congenital hypothyroidism. Ann Clin Biochem 1981 Nov;18(Pt 6):361-3

- Low LC; Lin HJ; Cheung PT; Lee FT; Chu SY; Kwok TL; Bacon Shone J; Yeung CY. Screening for congenital hypothyroidism in Hong Kong. Aust Paediatr J 1986 Feb;22(1):53-6
- Lur'e IV; Nikolaev DL. [Congenital developmental defects in the genetic monitoring system. I. The selection of developmental defects subject to dynamic observation]. Genetika 1983;19(1):165-70 [Rus]
- Lyon IC. Screening for congenital hypothyroidism: a three year experience. NZ Mcd J 1984 Mar 28;97(752):175-7
- Lyon IC; Crossley JR; Smith PA. Screening for cystic fibrosis. NZ Med J 1983 Sep 14;96(739):673-5
- Lyon IC; Webster DR. Newborn screening for homocystinuria [letter]. NZ Med J 1986 Aug 13;99(807):600-1
- Lyon ICT; Mitchell EA; Atherton EK; Webster DR. Screening for biotinidase deficiency. N Engl J Med 1986;314(22):1457
- Lyon, Jeff. Playing God in the Nursery. New York: W.W. Norton, 1985.

M				

- Machill G; Knapp A. [Sensitivity of phenylketonuria screening].

 Kinderarztl Prax 1984 Jul;52(7):347-50 [Ger]
- Macris MP; Nahmias AJ; Bailey PD; Lee FK; Visintine AM; Brann AW. Electron microscopy in the routine screening of newborns with congenital cytomegalovirus infection. J Virol Methods 1981 Jun; 2(6):315-20
- Mahle M; Schmidt H; Lutz P; Wendel U. [Initiation of treatment following screening for phenylketonuria]. Monatsschr Kinderheilkd 1984 Nov; 132(11):865-6 [Ger]
- Malone JI. Infant screening for phenylketonuria in Florida. J Fla Med Assoc 1983 Sep;70(9):782-3
- Mamunes P. Neonatal screening tests. Pediatr Clin North Am 1980 Nov; 27(4):733-51
- Manca L; Formato M; Demuro P; Pilo G; Gallisai D; Orzalesi M; Masala B. The gamma globin chain heterogeneity of the Sardinian newborn baby. Hemoglobin 1986;10(5):519-28
- Manning DN. Dried blood spots: a convenient technique for neonatal hormone screening. Med Lab Sci 1982 Jul;39(3):257-60
- Marble RD; Thomas RG; Sterling ML. Screening for angel dust in newborns [letter]. Pediatrics 1980 Aug;66(2):334

- Marchi AG; Castelpietra L; Mangiarotti MA; Cigui I; Uxa F; Pecile V; Nordio S. [The BM test and lysosomal enzyme in the meconium in neonatal screening of cystic fibrosis]. Minerva Pediatr 1981 Jul 15; 33(13):641-6 [Ita]
- Mardesic D; Dumic M; Tajic M; Gjuric G; Letica Protega N. Estimation of 17 alpha-hydroxyprogesterone in dried blood spots for mass screening of newborn infants with congenital adrenal hyperplasia. Acta Med Iugosl 1983;37(2):95-100
- Mastella G; Barlocco EG. Neonatal screening for cystic fibrosis [letter]. Lancet 1986 May 17;1(8490):1162-3
- Mastella G; Borgo G; Castellani E; et al. [Albumin, alpha-1-antitrypsin and lactase activity assays in meconium for the neonatal screening of cystic fibrosis]. Riv Ital Pediatr 1981;7(5):581-91 [Ita]
- Mastella G; Pederzini F; Girella E; Righetti G; Zanchetta M; Rizzotti P. Cystic fibrosis screening policy [letter]. Lancet 1984 Sep 8;2(8402):575-6
- Matalon R; Michals K; Lee CL; Nixon JC. Screening for biopterin defects in newborns with phenylketonuria and other hyperphenylalaninemias. Ann Clin Lab Sci 1982 Sep-Oct;12(5):411-4
- Mathias D. [Experience from 17 years of neonatal screening for inborn errors of metabolism in the Federal Republic of Germany, 1969-1985]. Arztl Lab 1986;32(11):263-6 [Ger]
- Mathias D; Bickel H. Follow-up study of 16 years neonatal screening for inborn errors of metabolism in West Germany. Eur J Pediatr 1986 Sep; 145(4):310-2
- Mathias D; Bickel H. Inter-laboratory quality control in neonatal screening for inborn errors of metabolism. Eur J Pediatr 1985 Jul; 144(2):171-3
- Matschke RG; Plath P. [Early detection of hearing disorders. A simple method of mass screening in newborn infants]. HNO 1985 Jan;33(1):40-4 [Gcr]
- Matsuda I; Matsuo K; Endo F; Uehara I; Nagata N; Jinno Y; Chikazawa S; Miyakita T; Miura H. Skin histidase activity and urine formiminoglutamic acid (FIGLU) in patients with histidinemia found by screening newborn infants. Clin Chim Acta 1982 Mar 12;119(3):319-28
- Matsumoto S; Miyamae T; Mashita M; et al. [An infant with ectopic thyroid glands without hypothyroidism found in neonatal mass screening]. J Saitama Med School 1981;7(4):393-7 [Jpn]
- Mayer HO; Stunzner D; Rosanelli K. [Value of a toxoplasmosis screening].
 Zentralbl Gynakol 1983;105(17):1097-100 [Ger]

- McCabe ER; McCabe L. Screening for PKU in sick or premature neonates [letter]. J Pediatr 1983 Sep;103(3):502-3
- McCabe ER; McCabe L; Mosher GA; Allen RJ; Berman JL. Newborn screening for phenylketonuria: predictive validity as a function of age. Pediatrics 1983 Sep;72(3):390-8
- McCormick B; Curnock DA; Spavins F. Auditory screening of special care neonates using the auditory response cradle. Arch Dis Child 1984 Dec;59(12):1168-72
- McCurdy PR. Screening for abnormal hemoglobins: who, when, and how. Primary Care 1980 Sep;7(3):451-60
- McCurdy RS. Newborn screening for genetic/metabolic diseases in Colorado. Colo Med 1983 Feb;80(2):48-50
- McFarland WH; Simmons FB; Jones FR. An automated hearing screening technique for newborns. J Speech Hear Disord 1980 Nov;45(4):495-503
- McGuinness GA; Smith WL. Head ultrasound screening in premature neonates weighing more than 1,500 g at birth. Am J Dis Child 1984 Sep; 138(9):817-20
- McNeil TF; Thelin T; Aspegren Jansson E; Sveger Γ; Harty B.

 Psychological factors in cost-benefit analysis of somatic prevention.

 A study of the psychological effects of neonatal screening for alpha 1-antitrypsin deficiency. Acta Paediatr Scand 1985 May;74(3):427-32
- McNeil TF; Thelin T; Aspegren JE; Sveger T. Identifying children at high somatic risk: possible effects on the parents' views of the child's health and parents' relationship to the pediatric health services.

 Acta Psychiatr Scand 1985 Dec;72(6):491-7
- McNeil TF; Thelin T; Aspegren JE; Sveger T. Identifying children at high somatic risk: possible long-term effects on parental conceptualizations of the children's personality and behavior. Acta Psychiatr Scand 1985 Dec;72(6):498-504
- Mehes K. Minor malformations in the neonate: utility in screening infants at risk of hidden major defects. Prog Clin Biol Res 1985;163C:45-9
- Mehes K; Juhasz E; Ruszinko V. Chromatographic screening of 70,328 neonates for inborn errors of amino acid metabolism. Acta Paediatr Hung 1985;26(2):147-9
- Meijer WJ. [Screening for congenital hypothyroidism]. Tijdschr Kindergeneeskd 1985 Jun;53(3):92-8 [Dut]
- Meijer WJ. [10 years' national screening study of the incidence of phenylketonuria in the Netherlands; 3d report of the National Advisory Phenylketonuria Commission]. Ned Tijdschr Geneeskd 1985 Jan 12; 129(2):74-6 [Dut]

- Melkova J; Pokorny A. [Detection rate of inborn heart defects in the newborn]. Cesk Pediatr 1981;36(5):254-7 [Cze]
- Mellis CM. Cystic fibrosis. Med J Aust 1985;143(6):227-8
- Meloni T; Gallisai D; Dore A; Forteleoni G; Mela G. Neonatal screening for hemoglobinopathy in North Sardinia. Eur J Pediatr 1981 Oct; 137(2):195-6
- Mengreli C; Kassiou K; Tsagaraki S; Pantelakis S. Neonatal screening for hypothyroidism in Greece. Eur J Pediatr 1981 Oct;137(2):185-7
- Merault G; Keclard L; Saint, Martin C; Jasmin K; Campier A; Delanoe, Garin J; Arous N; Fortune R; Theodore M; Seytor S; et al. Hemoglobin Roseau-Pointe a Pitre alpha 2 beta 2(90) (F6) Glu----Gly: a new hemoglobin variant with slight instability and low oxygen affinity. FEBS Lett 1985 May 6;184(1):10-3
- Meryash DL; Levy HL; Guthrie R; Warner R; Bloom S; Carr JR.

 Prospective study of early neonatal screening for phenylketonuria. N

 Engl J Med 1981 Jan 29;304(5):294-6
- Milunsky, Aubrey; Annas, George J. Genetics and the Law III. Proceedings of the Third National Symposium on Genetics and the Law; cosponsored by the American Society of Law and Medicine and the Boston Univ. Schools of Medicine, Law, and Public Health; Boston, 2-4 Apr 1984. New York: Plenum Press, 1985. 514 p.
- Misumi H; Wada H; Kawakami M; Ninomiya H; Sueishi T; Ichiba Y; Shohmori T. Detection of UDP-galactose-4-epimerase deficiency in a galactosemia screening program. Clin Chim Acta 1981 Oct 8;116(1): 101-5
- Mitchell ML; Bapat V; Larsen PR; et al. Pitfalls in screening for neonatal hypothyroidism. Pediatrics 1982;70(1):16-20
- Miyai K. [Neonatal screening--present and future]. Rinsho Byori 1986 Jun; 34(6):658-65 [Jpn]
- Miyai K; Ishibashi K; Kawashima M. Enzyme immunoassay for measuring thyrotropin in dried blood samples on filter paper for neonatal hypothyroid screening. J Clin Chem Clin Biochem 1981;19(8):774
- Miyai K; Tsuruhara T; Kusuda S; et al. Semiautomated enzyme immunoassay of thyrotropin as a mass screening test for neonatal hypothyroidism. Pediatr Res 1984;18(12):1289-92
- Miyaji T; Matsumoto S. [An infant with TBG (thyroxine binding globulin) deficiency and his family found during newborn thyroid screening]. J Saitama Med School 1981;8(3):157-62 [Jpn]

- Miyao M; Kuroda Y. Tentative classification of histidinemia detected by screening newborn infants on the as of skin histidase activity. Acta Paediatr Jpn 1982;24(1):147-50
- Mizuno T; Abe N; Teshima H; Yamauchi E; Itagaki Y; Matsumoto I; Kuhara T; Shinka T. Application of a gas chromatography mass spectrometry computer system for clinical diagnosis. Biomed Mass Spectrom 1981 Dec;8(12):593-7
- Mizushima Y; Fukushi M; Arai O; Sato T; Sato Y; Hayashi H. [The evaluation of mass-screening for congenital hypothyroidism by enzyme immunoassay for thyroid stimulating hormone]. Horumon To Rinsho 1983 Jan;31(1):69-76 [Jpn]
- Mizuta H. [III. Clinical diagnosis by immunoassay: mass screening]. Rinsho Byori 1983 Feb;Suppl 53:134-46 [Jpn]
- Mizuta P. Miyai K; Ichihara K; Amino N; Harada T; Nose O; Tanizawa O. Radioimmunoassay of "free thyroxin" in dried blood spots on filter paper preliminary observations on the effective differentiation of subjects with congenital hypothyroidism from those with subnormal thyroxin-binding globulin and normal subjects. Clin Chem 1982 Mar; 28(3):505-8
- Monk CJE; Dowed GSE. Monthly screening in the first six months of life for congenital hip dislocation. Isr J Med Sci 1980;16(4):253-6
- Monplaisir N; Casius de Linval JC; Sellaye M; Galacteros F; Braconnier F; Beuzard Y; Hilbert J; Mezin R; Quist C; Duville S; Rosa J. [Detection of hacmoglobinopathies at birth, using isoelectric focalization (author's transl)]. Nouv Presse Med 1981 Oct 24;10(38): 3127-30 [Fre]
- Moore H; McMillan M. Development of a filter paper blood spot radioimmunoassay for thyroid stimulating hormone suitable for a regional neonatal screening unit. Ann Clin Biochem 1983 Mar;20 Pt 2: 93-9
- Moosa A. Duchenne's muscular dystrophy in six siblings. The case for early diagnosis and neonatal screening. S Afr Med J 1982 Nov 13;62(21):765-7
- Morales WJ; Lim DV; Walsh AF. Prevention of neonatal group B streptococcal sepsis by the use of a rapid screening test and selective intrapartum chemoprophylaxis. Am J Obstet Gynecol 1986 Nov;155(5):979-83
- More on newborn screening for phenylketonuria: recommendations of the Committee on Genetics [letter]. Pediatrics 1983 Jul;72(1):139-42
- Morissette J; Dussault JH. Natural evolution of thyrotropin and thyroid hormones concentrations in hypothyroid newborns with thyroid agencsis or ectopy during the first month of life. Clin Invest Med 1986;9(3): 198-200

- Morris AF; Holton JB; Burman D; Colley JRT. Phenylalanine and tyrosine levels in newborn screening blood samples. Arch Dis Child 1983; 58(4):271-5
- Moschini L; Natoli G; Boscherini B; et al. [Congenital hypothyroidism. Two years' experience in neonatal screening]. Riv Ital Pediatr 1980; 6(6):693-9 [Ita]
- Moser H. [Genetic family counseling. Ethical viewpoints]. ZFA (Stuttgart) 1983 Jun 20;59(17):967-78 [Ger]
- Mudd SH; Skovby F; Levy HL; et al. The natural history of homocystinura due to cystathionine beta-synthase deficiency. Am J Hum Genet 1985; 37(1):1-31
- Murata M; Nakajima K; Matsumoto M. [Some problems on the screening of neonatal cretinism (author's transl)]. Horumon To Rinsho 1981 Mar;29(3):289-92 [Jpn]
- Murata T. [Studies on judgement of high risk neonates. II. New screening system for the estimation of neonatal mortality and identifying of the high risk neonate]. Acta Neonatal Jpn 1981;17(3):461-8 [Jpn]
- Murray AD; Javel E; Watson CS. Prognostic validity of auditory brainstem evoked response screening in newborn infants. Am J Otolaryngol 1985 Mar-Apr;6(2):120-31

N					
			 		_

- Nadler HL. Urine screening after discharge from hospital. Pediatrics 1981; 67(1):158-9
- Nagataki S; Ishibashi K; Ohsawa R; et al. Mcasuring thyroxinc and thyrotropin simultaneously in a dried blood sample on filter paper, to screen for neonatal hypothyroidism. Clin Chem 1980;26(6):750-3
- Naruse H; Ishii S; Momose T; et al. Scnsitive EIA methods for determining thyrotropin and thyroxine in dried blood on filter paper: New methods for neonatal thyroid screening. J Clin Chem Clin Biochem 1981;19(8):781
- Naruse H; Suzuki E; Irie M; Tsuji A; Takasugi N; Fukushi M; Matsuura N; Shimozawa K. Neonatal screening for congenital adrenal hyperplasia in Japan. Ann NY Acad Sci 1985;458:103-10
- National Clearinghouse for Human Genetic Diseases. State Laws and Regulations on Genetic Disorders. Rockville, Md.: Bureau of Community Health Services, 1980. 80 p.
- Naylor EW. Newborn screening for urea cycle disorders. Adv Exp Med Biol 1982;153:9-18

- Naylor EW. Newborn screening of urea cycle disorders. Pediatrics 1981 Sep; 68(3):453-7
- Naylor EW. Recent developments in neonatal screening. Semin Perinatol 1985 Apr;9(3):232-49
- Nekvasil R; Kucerova I; Kalinova H. [Screening for anemia and polycythemia in the early neonatal period]. Cesk Pediatr 1981 Apr; 36(4):230-1 [Cze]
- Nekvasil R; Stejskal J; Tuma A. Detection of early onset neonatal hypocalcaemia in low birth weight infants by Q-Tc and Q-oTc interval measurement. Acta Paediatr Acad Sci Hung 1980;21(4):203-10
- Nelson MF. Wrongful life: impaired infant's cause of action recognized--«Curlender v. Bio-Science Laboratories». Brigham Young Univ Law Rev 1980;1980(3):676-683
- Neonatal hypothyroidism screening: Status of patients at 6 years of age. J Pediatr 1985;107(6):915-9
- Neonatal screening for cystic fibrosis [letter]. Lancet 1986 Feb 8; 1(8476):326-8
- Neonatal screening for cystic fibrosis [letter]. Lancet 1986 Mar 15; 1(8481):615
- Neonatal screening for cystic fibrosis: position paper. Pediatrics 1983 Nov;72(5):741-5
- Neonatal screening for phenylketonuria [letter]. Pediatrics 1984 May; 73(5):737-40
- Netzloff ML. Neonatal screening for phenylketonuria. Ann Clin Lab Sci 1982 Sep-Oct;12(5):368-71
- Newborn screening. District of Columbia Code Annotated. 1981. Sections 6-311 to 6-320, added 1980. 5 p.
- Newborn screening for metabolic disorders in Australia and New Zealand: results for 1983. Human Genetics Society of Australasia Newborn Screening Committee. Med J Aust 1985 Aug 19;143(4):159-60
- Nielsen J; Wohlert M; Faaborg Andersen J; et al. Incidence of chromosome abnormalities in newborn children. Comparison between incidences in 1969-1974 and 1980-1982 in the same area. Hum Genet 1982;61(2):98-101
- Nordshus T; Monn E. Ultrasonography in the diagnosis of neonatal adrenal haemorrhage. Acta Paediatr Scand 1980 Sep;69(5):695-8
- Nose O; Harada T; Kodama H; Kai H; Ogawa M. [Diagnostic procedure of neonatal hypothyroidism detected by a mass screening program (author's transl)]. Horumon To Rinsho 1981 Mar;29(3):313-8 [Jpn]

- Novogroder M. Neonatal screening for congenital hypothyroidism. Pediatr Clin North Am 1980 Nov;27(4):881-90
- Nowak W; Danielewicz B; Mika Bortkiewicz M. [Uscfulness of screening tests in neonatal diagnostics]. Pediatr Pol 1980 Mar;55(3):369-71 [Pol]
- Nussbaum RL; Powell C; Graham HL; Caskey CT; Fernbach DJ. Newborn screening for sickling hemoglobinopathies. Houston, 1976 to 1980. Am J Dis Child 1984 Jan;138(1):44-8
- Nyhan WL. Neonatal screening for inherited disease [editorial]. N Engl J Med 1985 Jul 4;313(1):43-44

^	^			
v	O .			

- Oberg CN; Sklar C. Congenital hypothyroidism. An imperfection of newborn screening. Minn Med 1985 Dec;68(12):889-90
- O'Halloran ET; Crowley MJ. Screening for cystic fibrosis. Ir Med J 1982 Dec;75(12):480-1
- Orfanos AP; Jinks DC; Guthrie R. Microassay for estimation of galactose and galactose-1-phosphate in dried blood specimens. Clin Biochem 1986 Aug;19(4):225-8
- Oura T. Genetic screening for metabolic disorders: Present state and issue.

 Japanese system of the newborn screening for inborn errors of metabolism. Jpn J Hum Genet 1986;31(2):121-2
- Oura T. [Present status and new problems in the newborn screening for inborn errors of metabolism]. Congen Anomal 1981;21(1):9-16 [Jpn]
- Outcome of screening for congenital hypothyroidism [editorial]. Lancet 1986 May 17;1(8490):1130-1

- Paglietti E; Galanello R; Moi P; Pirastu M; Cao A. Molecular pathology of haemoglobin H disease in Sardinians. Br J Haematol 1986 Jul; 63(3):485-96
- Paigen K; Pacholec F; Levy HL. A new method of screening for inherited disorders of galactose metabolism. J Lab Clin Med 1982;99(6):895-907
- Pandav CS; Kochupillai N. Organisation and implementation of neonatal hypothyroid screening programme in India--a primary health care approach. Indian J Pediatr 1985 May-Jun;52(416):223-9



- Pang S; Murphey W; Levine LS; Spence DA; Leon A; LaFranchi S; Surve AS; New MI. A pilot newborn screening for congenital adrenal hyperplasia in Alaska. J Clin Endocrinol Metab 1982 Sep;55(3):413-20
- Pang S; Spence DA; New MI. Newborn screening for congenital adrenal hyperplasia with special reference to screening in Alaska. Ann NY Acad Sci 1985;458:90-102
- Pantelakis S; Lambadaridis I; Mengreli C. Forty eight cases of primary congenital hypothyroidism diagnosed with the neonatal screening programme. Acta Endocrinol [Suppl] 1983;104(261):21-2
- Park JM; Bourgeois MJ; Lohla E; Varma SK. Cardiac function of neonates with congenital hypothyroidism before treatment. Am J Med Sci 1986 Sep; 292(3):157-9
- Parkin DM. How successful is screening for congenital disease of the hip?

 Am J Public Health 1981 Dec;71(12):1378-83
- Passos MR; Zatz M. Creatine-kinase (CK) and pyruvate-kinase (PK) activities in cord blood of normal newborn infants: application to Duchenne muscular dystrophy screening programs. Am J Med Genet 1983 Nov;16(3):367-72
- Paul TD; Naylor EW; Guthrie R. Urine screening for metabolic disease in newborn infants. J Pediatr 1980 Apr;96(4):653-6
- Perrine RP; John P; Pembrey M; Perrine S. Sickle cell disease in Saudi Arabs in early childhood. Arch Dis Child 1981 Mar; 56(3):187-92
- Petersen PH; Rosleff F; Rasmussen J; Hobolth N. Studies on the required analytical quality of TSH measurements in screening for congenital hypothyroidism. Scand J Clin Lab Invest [Suppl] 1980;155:85-93
- Phadke MA; Metelu S; Kate SL; Joshi VS; Shirole DB; Joshi AS; Mokashi GD. Screening of newborn for inborn errors of metabolism. Indian Pediatr 1982 Sep;19(9):767-9
- Phenylketonuria. Illinois Annotated Statutes (Smith-Hurd). 1982. Chapter 111 1/2, Sections 4901 to 4905. 2 p.
- Philip AG. Decreased use of antibiotics using a neonatal sepsis screening technique. J Pediatr 1981 May;98(5):795-9
- Philip AG. Detection of neonatal sepsis of late onset. JAMA 1982 Jan 22-29;247(4):489-92
- Phuapradit P; Narang A; Mendonca P; et al. The steatocrit: A simple method for estimating stool fat content in newborn infants. Arch Dis Child 1981;56(9):725-7

- Piazzi S; Capelli M; Martelli E; Paolini M; Tolomelli M; Cassio A;

 Balsamo A; Bugiardini G. Problems connected with neonatal screening assays. J Nucl Med Allied Sci 1982 Oct-Dec;26(4):319-25
- Piazzi S; Capelli M; Paolini M; Perugini D; Grossi G; Balsamo A; Salomoni P; Cassio A; Bugiardini G; Cacciari E. Neonatal screening for 21-hydroxylase deficiency: a microfilter paper method for 17-alpha-hydroxyprogesterone assay. J Endocrinol Invest 1982 Mar-Apr; 5(2):87-90
- Piccolo L; Botner F. [Newborn hearing screening in the hospital of Mestre from Nov. '72 to Dec. '78]. Acta Phoniatr Lat 1980;2(1):57-60 [Ita]
- Pierson EA; Therrell BL Jr; Franz SF; et al. An evaluation of kits for measuring thyrotropin in newborns. Clin Chem 1984;30(11):1854-6
- Pitfalls in screening for neonatal hypothyroidism. Report of the New England Regional Screening Program and the New England Congenital Hypothyroidism Collaborative. Pediatrics 1982 Jul;70(1):16-20
- Pitt D; Connelly J; Francis I; Wilcken B; Brown DA; Robertson E; Hill G; Masters P; Tucker RG; Raby J; McFarlane J; Hancock J. Genetic screening of newborn in Australia: results for 1980. Mcd J Aust 1982 Feb 6;1(3):119-20
- Pitt D; Connelly J; Francis I; Wilchen B; Brown DA; Robertson E; Hill G; Masters P; Raby J; McFarlane J; Bowling F; Hancock J. Genetic screening of newborn in Australia. Results for 1981. Med J Aust 1983 Apr 2;1(7):333-5
- Pitt D; Connelly J; Francis I; Wilcken B; Brown DA; Robertson E; Hill G; Masters P; Tucker RG; Raby J; McFarlane J; Hancock J. Genetic screening of newborn in Australia: results for 1979. Mcd J Aust 1981 Jan 10;1(1):39-40
- Plauchu H; Dellamonica C; Cotte J; Robert JM. Duchenne muscular dystrophy: systematic neonatal screening and earlier detection of carriers. J Genet Hum 1980 Dec;28(4):65-82
- Plauchu H; Dellamonica C; Cotte J; Robert JM. [Noonatal screening for Duchenne muscular dystrophy. The Lyons experience]. Rev Med Liege 1984 May 15;39(10):464-8 [Frc]
- Plauchu H; Dellamonica C; Pascal B; Jolivet MJ; Guibaud P; Cotte J; Robert JM. [Nconatal screening for duchenne myopathy by scrum elevation of creatine phosphokinase activity. 5 years experience]. J Genet Hum 1981 Mar;29(1):59-69 [Fre]
- Pollitt RJ. Neonatal screening in France. J Inherited Metab Dis 1983;6(2):49-50

- Pollitt RJ; Worthy E; Green A. Galactosaemia detection as a bonus from screening for phenylketonuria. J Inherit Metab Dis 1982;5(Suppl.1):51-2
- Pomarede R; Czernichow P; Farriaux JP. [Analysis of failures of screening for congenital hypothyroidism by the assay of TSH in capillary blood]. Arch Fr Pediatr 1986 Jan;43(1):15-8 [Fre]
- Popescu A; Butnariu J; Toma C; Farcas A; Gelmareanu K. [Data on phenylketonuria screening]. Rev Pediatr Obstet Ginecol [Pediatr] 1983 Jan-Mar; 32(1):67-9 [Rum]
- Porter IH. Control of hereditary disorders. Annu Rev Public Health 1982; 3:277-319
- Postellon DC. Detection of congenital nephrosis by screening program for hypothyroidism [letter]. J Pediatr 1983 Feb;102(2):321-2
- Postellon DC. Diagnosis and treatment of congenital hypothyroidism. Compr Ther 1983;9(2):41-4
- Postellon DC. Screening for congenital hypothyroidism [letter]. J Pediatr 1981 Jul;99(1):170-1
- Poyhonen L; Lenko HL. Ultrasonography in congenital hypothyrcosis. Acta Fuediatr Scand 1984 Jul;73(4):523-6
- Prabt kar P; Singhi S; Sharma A; James O. Immunoglobulin and C3 levels ramaica. Trop Geogr Med 1985 Dec;37(4):304-8
- Pr. S; Reese L; Tevaarwerk GJM; et al. Relation of cord blood roxine and thyrotropin levels to gestational age and birth weight.

 (1 Med Assoc J 1980;123(10):1007-13
- Prekop R; Sedlak J; Varady L; Pillarova L. [Results of screening for congenital hypothyroidism in hypotrophic neonates]. Cesk Pediatr 1985 Oct;40(10):573-5 [Slo]
- Price DA; Ehrlich RM; Walfish PG. Congenital hypothyroidism. Clinical and laboratory characteristics in infants detected by neonatal screening. Arch Dis Child 1981;56(11):845-51
- Primhak RA; Simpson RM. Screening small for gestational age babies for congenital infection. Clin Pediatr (Phila) 1982 Jul;21(7):417-20

R

Radu HJ. [Comparison of different audiometric procedures for infants less than 1 year old]. HNO 1985 Jun;33(6):271-4 [Ger]

- Ralston KK; Kmetz DR; Keeling MM; Queenau JT. Screening for major hemoglobinopathics in newborn Blacks. J Ky Mcd Assoc 1981 Oct;79(10): 649-51
- Ramamurthy RS; Brans YW. Neonatal polycythemia: I. Criteria for diagnosis and treatment. Pediatrics 1981 Aug;68(2):168-74
- Rand KH; Houck HJ; Swingle HM. Rotazyme assay in neonates without diarrhea. Results of screening survey and preliminary analysis of false positive specimens. Am J Clin Pathol 1985 Dec;84(6):748-51
- Rao S; Thurston AJ. Congenital dislocation of hip in the newborn: a postnatal survey. NZ Med J 1986 Oct 8;99(811):752-4
- Rappaport R. [Myxedema in children]. Concours Med 1981;103(23):3857-60 [Fre]
- Reddy UB. Delaware's hereditary metabolic disorders infant screening program. Del Med J 1982 Oct;54(10):555-7
- Richmond KH; Konkle DF; Potsic WP. ABR screening of high-risk infants: effects of ambient noise in the neonatal nursery. Otolaryngol Head Neck Surg 1986 Jun;94(5):552-60
- Rigillo N; Zaccaro A; Pisano R; Trisolini S. [Neonatal screening program for cystic fibrosis of the pancreas]. Pediatria (Napoli) 1981 Dec 31; 89(4):747-60 [Ita]
- Rivera de Caragol E. [Neonatal screening: who is covered? (letter)] Bol Asoc Med PR 1986 Sep;78(9):411 [Spa]
- Roberton NR. Screening for congenital hip dislocation [letter]. Lancet 1984 Apr 21;1(8382):909-10
- Robertson EF; Wilkins AC; Oldfield RK; Pollard AC. Blood spot thyroxine-binding globulin: a means to reduce recall rate in a screening strategy for neonatal hypothyroidism. J Pediatr 1980 Oct;97(4):604-7
- Robinson A. Genetic screening: medical progress prompts ethical questions. AORN J 1986 May;43(5):1137, 1140, 1142
- Rochiccioli P; Dutau G. [Hypothyroidism in infants and children]. Rev Pediatr 1981;17(1):21-6 [Fre]
- Rochiccioli P; Dutau G; Augier D. [Thyroid ectopia, a cause of error in neonatal screening for hypothyroidism]. Arch Fr Pediatr 1983 May; 40(5):405-6 [Fre]
- Rochiccioli P; Dutau G; Enjaume C; et al. [Screening for neonatal hypothyroidism by measurement of both T4 and TSH in dried blood eluates]. Arch Fr Pediatr 1980;37(3):159-62 [Fre]

i.

- Rochiccioli P; Dutau G; Roge B; Petrus M; Augier D; Enjaume C. [Neonatal screening of hypothyroidism in France. One million quantitative analyses and psycho-motor development of 20 detected cases]. J Genet Hum 1981 Mar;29(1):13-22 [Fre]
- Roesel RA; Blankenship PR; Hommes FA. HPLC assay of phenylalanine and tyrosine in blood spots on filter paper. Clin Chim Acta 1986 Apr 15; 156(1):91-6
- Romagnoli C; Curro V; Luciano R; Tortorolo G; Segni G; Carta Sorcini M; Tomarchio S; Fiore L; Di Iorio MG; Gilardi E; Carta S. Scrial blood T4 and TSH determinations in low birth weight infants, influence of gestational age, birth weight and neonatal pathology on thyroid function. Helv Paediatr Acta 1982 Sep;37(4):331-44
- Romagnoli C; Curro V; Luciano R; Zecca E; De Carolis MP; Tortorolo G; Carta Sorcini M; Tomarchio S; Alessandrini L; Fiore L; et al. [Hypothyroxinemia in the low birth-weight infant in the screening of congenital hypothyroidism]. Pediatr Mcd Chir 1982 Nov-Dec;4(6):679-84 [Ita]
- Romeo G; Menozzi P; Ferlini A; Prosperi L; Cerone R; Scalisi S; Romano C; Antonozzi I; Riva E; Piceni Sereni L; et al. Incidence of classic PKU in Italy estimated from consanguineous marriages and from neonatal screening. Clin Genet 1983 Nov;24(5):339-45
- Ronconi G; Tezzele A; De Boni P; Fantuz E; Toniolo S. [Screening for neonatal hypothyroidism by estimation of T4 and TSH. Experience of three years on 6.340 new-borns (author's transl)]. Pediatr Med Chir 1981 Mar-Jun;3(2-3):199-202 [Ita]
- Ronconi GF; Pesenti P; Tezzele A; Bogoni A; Zanardo V. [Two cases of familial thyroxine-binding globulin (TBG) deficit identified by the screening of neonatal hypothyroidism (author's transl)]. Pediatr Med Chir 1981 Mar-Jun;3(2-3):239-42 [Ita]
- Root AW. Current status of neonatal screening for congenital hypothyroidism. Compr Ther 1986;12(2):51-4
- Rosati U; Bigo E; Picco P; et al. [Neonatal screening for congenital hypothyroidism in Regione Liguria]. G Ig Med Prev 1981;22(1):25-30 [Ita]
- Rosati U; Minicucci L; Cotellessa M; et al. [Program of newborn screening for genetic diseases. Theories and operative basis]. G Ig Med Prev 1981;22(1):7-12 [Ita]
- Rosati U; Scalisi S; Minicucci L; et al. [Neonatal screening for PKU in Regione Liguria]. G Ig Med Prev 1981;22(1):13-17 [Ita]
- Rosenthal IM. Screening of the newborn for hypothyroidism. Indian J Pediatr 1982 Mar-Apr;49(397):273-6

- Roth KS. Newborn metabolic screening: a search for "nature's experiments." South Med J 1986 Jan;79(1):47-54
- [Round table: neonatal screening programs in Europe]. J Genet Hum 1981 Mar;29(1):23-38 [Frc]
- Routine neonatal screening for phenylketonuria in the United Kingdom 1964-78. Medical Research Council Steering Committee for the MRC/DHSS Phenylketonuria Register. Br Med J [Clin Res] 1981 May 23; 282(6277):1680-4
- Rovet JF. A prospective investigation of children with congenital hypothyroidism identified by neonatal thyroid screening in Ontario. Can J Public Health 1986 May-Jun;77 Suppl 1:164-73
- Rowley PT. Genetic screening: marvel or menace? Science 1984 Jul 13; 225(4658):138-44
- Rowley PT; Huntzinger DJ. Newborn sickle cell screening. Benefits and burdens realized. Am J Dis Child 1983 Apr;137(4):341-5
- Rucknagel DL. A decade of screening in the hemoglobinopathies: is a national program to prevent sickle cell anemia possible? Am J Pediatr Hematol Oncol 1983 Winter;5(4):373-7
- Rudolph WC. Social issues in perinatal screening for sickle cell anemia. Urban Health 1983 Aug;12(8):32-4
- Ruhtenberg M; Wagenknecht C; Miller EC. [Amino acid imbalances in blood of pregnant women with diabetes and in blood of their newborns--limit values for screening (author's transl)]. Zentralbl Gynakol 1982;104(9):519-25 [Ger]
- Ruhtenberg M; Wagenknecht C; Miller EC; Kohlhoff R. [Amino acid imbalance in the blood of pregnant diabetic women and their newborn infants--limit values for screening. II. Neonatal amino acid imbalance]. Zentralbl Gynakol 1982;104(16):1028-37 [Ger]
- Russo V. Quality assurance: Improving neonatal screening for inherited metabolic diseases. J Am Med Rec Assoc 1983;54(4):24-9
- Rybakowa M; Markowska Kraik E; Ostrowska J; Sotysik Wilk E; Sztefko K; Tomaszczyk B. [TSH and thyroid hormone levels in newborn infants with hyperthyrotropinemia detected by screening tests, in newborn infants with goiter and those born to mothers with hyperthyroidism]. Endokrynol Pol 1985;36(5):277-84 [Pol]

S

Sack J; Amado O; Frucht H; Ekstein A. Screening for neonatal hypothyroidism in Israel. Isr J Med Sci 1981 Apr;17(4):294-5

- Sack J; Elicer A; Sofrin R; Theodor R; Cohen B. Influence on psychological development of early treatment of congenital hypothyroidism detected by neonatal screening: a controlled study. Isr J Med Sci 1986 Jan;22(1):24-8
- Sack J; Kletter G; Amado O; Akstein E. Screening for neonatal hypothyroidism in Israel during a 4-year period. Isr J Med Sci 1985 Jun;21(6):485-9
- Sahota A; Blair JA; Barford PA; Leeming RJ; Green A; Pollitt RJ.
 Neonatal screening for dihydropteridine reductase deficiency. J
 Inherited Metab Dis 1985;8 Suppl 2:99-100
- Saito T; Satoh K; Yajima Y; Nagai F. [Thyroxin in dried blood spot as a primary index in mass screening program for neonatal hypothyroidism (author's transl)]. Kaku Igaku 1980 Jul;17(6):747-54 [Jpn]
- Sakaguchi M; Suzuki S; Nagashima K; Shimano S; Uchida S; Kuroume T. Longitudinal study of free thyroxine in low-birth-weight infants by paper disk method. J Pediatr 1983 Nov;103(5):793-6
- Sakaida H; Imanishi H; Kitamura T. [Predictive value of acute phase reactants (APR)-score for perinatal infections]. Nippon Sanka Fujinka Gakkai Zasshi 1983 Jul;35(7):929-33 [Jpn]
- Sander J; Niehaus C. [Congenital hypothyroidism: results of screening 30,000 neonates in lower saxony (author's transl)]. MMW 1980 Feb 29; 122(9):309-12 [Ger]
- Sander J; Niehaus C. [Determination of free thyroxine (FT4) for the detection of connatal hypothyroidism within the scope of neonatal screening]. Monatsschr Kinderheilkd 1984 Nov;132(11):861-4 [Ger]
- Sander J; Niehaus C. [Hepatitis A antibodies in young women in Lower Saxony. Results of a serologic study on dried blood samples from the neonatal screening program for inborn errors of metabolism]. Off Gesundheitswes 1982 Apr;44(4):235-6 [Ger]
- Sander J; Niehaus C. [Mucoviscidosis screening by determination of immunoreactive trypsin]. Klin Padiatr 1984 Jul-Aug;196(4):224-7 [Ger]
- Sander J; Niehaus C. [Radio-immuno-assay for trypsin in newborn-screening for cystic fibrosis]. Monatsschr Kinderheilkd 1982 Nov;130(11):843-5 [Ger]
- Sander J'; Niehaus C. Screening test for syphilis-specific antibodies added to a screening program for congenital hypothyroidism. J Pediatr 1982 Jan;100(1):93-5
- Sander J; Niehaus Ch. [Rubella screening using the haemolysis-in-gel test from dried newborn blood on filter paper]. Dtsch Med Wochenschr 1980; 105(23):827-9 [Ger]

- Sander J; Niehaus Ch. [Screening for syphilis: results of a study of 25 000 neonates in Lower Saxony]. Dtsch Med Wochenschr 1980;105(24):858-60 [Ger]
- Sander J; Niehaus Ch. [Simultaneous thyroxin determination in thyrotropin (TSH) screening for congenital hypothyroidism]. Dtsch Med Wochenschr 1982;107(4):138-42 [Ger]
- Santiago Borrero PJ; Caceres de Costas M; Valcarcel M. Abnormal hemoglobins detected in screening of Puerto Rican population. Bol Asoc Med PR 1981 Jun;73(6):274-8
- Sardet A. [Allergic diseases: Screening and prevention in neonates and infants]. Arch Fr Pediatr 1983;40(10):809-14 [Fre]
- Sasaki H. Development of bile acid metabolism in neonates during perinatal period. Part 2: Mass screening of congenital biliary atresia by radioimmunoassay using dried blood spot. Acta Paediatr Jpn 1984; 26(2):161-8
- Sayeg N; Jumaa M; Sitzmann FG. Estimation of gene frequency of classical galactosemia in Syria. Ann Univ Sarav Med 1982;24(Suppl.2):23-6
- Schaub J. [Ethics in pediatric reseach]. Monatsschr Kinderheilkd 1982; 130(11):817-22 [Ger]
- Scheuerbrandt G. Screening for the early detection of Duchenne muscular dystrophy. Cardiomyology 1982;1(2):29-41
- Scheuerbrandt G; Lundin A; Lovgren T; Mortier W. Screening for Duchenne muscular dystrophy: an improved screening test for creatine kinase and its application in an infant screening program. Muscle Nerve 1986 Jan; 9(1):11-23
- Schmidt BJ; Diament AJ; Krynski S; et al. Neonatal mass screening of hereditary metabolic diseases in S. Paulo- Brasil. Acta Paediatr Jpn 1982;24(1):75-82
- Schmidt BJ; Diament AJ; Krynski S; Kamei ME; Rodrigues MM; Takata S. [Detection of inborn errors of metabolism in San Pablo, Brazil]. Bol Med Hosp Infant Mex 1981 Mar-Apr;38(2):217-29 [Spa]
- Schneider AJ. Newborn phenylalanine/tyrosine metabolism. Implications for screening for phenylketonuria. Am J Dis Child 1983;137(5):427-32
- Schoen EJ; Fefferman R. Screening for congenital hypothyroidism [letter].
 J Pediatr 1980 Jan;96(1):171-3
- Schonberg D; Klett M. [Neonatal screening for hypothyroidism; Experience in the GFR]. Therapiewoche 1980;30(42):6860-70 [Ger]
- Schreier K. [Modern possibilities to early diagnosis of congenital metabolic diseases]. Med Welt 1982;33(13):462-5 [Ger]

- Schwartz RP. Screening for congenital hypothyroidism in North Carolina-1979-1984. NC Med J 1985 Apr;46(4):205-7
- Scott RB; Harrison DL. Screening of the umbilical cord blood for sickle cell disease: utilization and implementation. Am J Pediatr Hematol Oncol 1982 Summer;4(2):202-5
- Screening and Counseling for Genetic Conditions: A Report on the Ethical, Social, and Legal Implications of Genetic Screening, Counseling, and Education Programs. President's Commission for the Study of Ethical Problems in Medicine and Biomedical and Behavioral Research. Washington, D.C.: The Commission, 1983. 122 p.
- Screening for congenital hypothyroidism [editorial]. Br Mcd J 1980 Jul 5;281(6232):1-2
- [Screening for congenital hypothyroidism]. S Afr Med J 1981;59(17):583
 [Afr]
- Screening for cystic fibrosis [letter]. J Pediatr 1983 Oct;103(4):667-8
- Screening for hearing impairment in the newborn [editorial]. Lancet 1986 Dec 20-27;2(8521-22):1429-30
- Screening for the detection of congenital dislocation of the hip. Arch Dis Child 1986 Scp;61(9):921-6
- Screening program for genetic/metabolic disorders. J Iowa Med Soc 1980 Feb;70(2):66, 69, 72
- Scriver CH. Genetic screening: implications for preventive medicine [editorial]. Am J Public Health 1983 Mar;73(3):243-5
- Scriver CR. Population screening: report of a workshop. Prog Clin Biol Res 1985;163B:89-152
- Scriver CR. Screening for medical intervention: the PKU experience. Prog Clin Biol Res 1982;103 Pt B:437-45
- Scriver CR; Holtzman NA; Howell RR; et al. New issues in newborn screening for phenylketonuria and congenital hypothyroidism. Pediatrics 1982;69(1):104-6
- Scriver CR; Levy HL. Histidinacmia. I. Reconciling retrospective and prospective findings. J Inherit Metab Dis 1983;6(2):51-3
- Sedlak J; Talan P; Prekop R. [RIA-Test-T 4 set (URVJT-Kosice) and its potential use in congenital hypothyroidism screening (author's transl)]. Cas Lek Cesk 1981 Feb 26;120(8):233-5 [Slo]
- Sengers RC. [Early detection of phenylketonuria in the Netherlands, 1977-1979. 2d report of the National Care Commission on Phenylketonuria]. Ned Tijdschr Geneeskd 1981 Dec 26;125(52):2135-40 [Dut]

- Sepe SJ. Screening for metabolic disease [letter]. J Pediatr 1980 Nov; 97(5):871-2
- Serjeant GR. Treatment of sickle cell disease in early childhood in Jamaica. Am J Pediatr Hematol Oncol 1985 Fall;7(3):235-9
- Serjeant GR; Serjeant BE; Forbes M; Hayes RJ; Higgs DR; Lehmann H. Haemoglobin gene frequencies in the Jamaican population: a study in 100,000 newborns. Br J Haematol 1986 Oct;64(2):253-62
- Shannon DA; Felix JK; Krumholz A; Goldstein PJ; Harris KC. Hearing screening of high-risk newborns with brainstem auditory evoked potentials: a follow-up study. Pediatrics 1984 Jan;73(1):22-6
- Sheftel DN; Hustead V; Friedman A. Hypertension screening in the followup of premature infants. Pediatrics 1983;71(5):763-6
- Shehab M; Barakat AY. Thalassemia B with distal renal tubular acidosis: a previously undescribed association. Int J Pediatr Nephrol 1985 Apr-Jun;6(2):143-4
- Shepard NT. Newborn hearing screening using the Linco-Bennett auditory response cradle: a pilot study. Ear Hear 1983 Jan-Feb;4(1):5-10
- Sheridan Pereira M; Gorman W. Transcutaneous bilirubinometry: An evaluation. Arch Dis Child 1982;57(9):708-10
- Shimozawa K. [Neonatal mass-screening for congenital adrenal hyperplasia due to 21-hydroxylase deficiency. II. A pilot neonatal mass-screening study in the west of Shizuoka Prefecture, Japan]. Nippon Naibunpi Gakkai Zasshi 1983 Dec 20;59(12):1860-73 [Jpn]
- Shimozawa K. [Neonatal mass-screening for congenital adrenal hyperplasia due to 21-hydroxylase deficiency. I. Microfilter paper methods for radioimmunoassay of 17 alpha-hydroxyprogesterone]. Nippon Naibunpi Gakkai Zasshi 1983 Dec 20;59(12):1845-9 [Jpn]
- Shimozawa K; Saisho S; Saito N; Yata J; Igarashi Y; Hikita Y; Irie M; Okada K. A neonatal mass-screening for congenital adrenal hyperplasia in Japan. Acta Endocrinol (Copenh) 1984 Dec;107(4):513-8
- Shuman BJ; Masterpasqua F. Preventive intervention during the perinatal and infancy periods: overview and guidelines for evaluation. Prev Hum Serv 1981 Fall-Winter;1(1-2):41-57
- Simmons FB. Patterns of deafness in newborns. Laryngoscope 1980 Mar; 90(3):448-53
- Singh H. Glucose-6-phosphate dehydrogenase deficiency: A preventable cause of mental retardation. Br Med J 1986;292(6517):397-8

- Skinner R; Emery AE; Scheuerbrandt G; Syme J. Fcasibility of neonatal screening for Duchenne muscular dystrophy. J Med Genet 1982 Feb; 19(1):1-3
- Skrabal J. [Results of dispensarization by three screenings when examining hip joints of the newborn (author's transl)]. Acta Chir Orthop Traumatol Cech 1980 Apr;47(2):143-53 [Cze]
- Snyderman SE; Sansaricq C. Newborn screening for maple syrup urine disease. J Pediatr 1985 Aug;107(2):259-61
- Solyom J; Hervei S; Marossy P; Solyom E; Babosa M; Szombathy G.
 "Dot-17 alpha-hydroxyprogesterone" radioimmunoassay for identification of congenital adrenal hyperplasia in young infants. Acta Paediatr Scand 1981 Nov;70(6):913-7
- Somers DG; Favreau L. Newborn screening for phenylketonuria: incidence and screening procedures in North America. Can J Public Health 1982 May-Jun;73(3):206-7
- Sood SK; Madan N; Talwar N; Maheshwari A; Jayant D; Bhargava SK. Reliability of three screening tests in detection of glucose-6-phosphate dehydrogenase deficiency in adults and neonates. Indian J Pathol Microbiol 1981 Apr;24(2):89-99
- Sorenson JR; Levy HL; Mangione TW; Sepe SJ. Parental response to repeat testing of infants with 'false-positive' results in a newborn screening program. Pediatrics 1984 Feb;73(2):183-7
- State Laws and Regulations on Genetic Disorders. Rockville, Md.: Bureau of Community Health Services, 1980. 80 p.
- Stein J; Berg C; Jones JA; Detter JC. A screening protocol for a prenatal population at risk for inherited hemoglobin disorders: results of its application to a group of Southeast Asians and blacks. Am J Obstet Gynecol 1984 Oct 15;150(4):333-41
- Stein L; Ozdamar O; Kraus N; Paton J. Follow-up of infants screened by auditory brainstem response in the neonatal intensive care unit. J Pediatr 1983 Sep;103(3):447-53
- Stempfel RS. Florida's infant screening program for metabolic disease. J Fla Med Assoc 1983 Sep;70(9):779-81
- Stine MJ. Newborn screening tests. Indiana Mcd 1986 Oct;79(10):868-9
- Stuhler T; Feige A; Kaesemann H. [The inset of sonography in completion to the orthopaedic hip-control at the newborn]. Z Orthop Ihre Grenzgeb 1985;123(2):121-6 [Ger]
- Sunartini Kodama S; Nishio H; et al. Infantile hypothyroidism discovered on neonatal screening in Hyogo prefecture Japan. ICMR Ann 1985;5:51-62

- Sutherland RM; Ratcliffe JG; Chapman RS. Immunoassay of blood spot TSH; development of a rapid two-site immunoradiometric assay and comparison with radioimmunoassay as a screening method for neonatal hypothyroidism. Clin Chim Acta 1982;124(1):1-11
- Sutherland RM; Ratcliffe JG; Kennedy R; Stevenson JS; Patrick MJ; Ferguson Smith MA. Neonatal screening for hypothyroidism in Scotland: results of a pilot study. Scott Med J 1981 Jul;26(3):229-34
- Suwa S; Yokoya S; Maesaka O; Sato M. [Infants with low-TBG detected in a screening program for neonatal hypothyroidism (author's transl)].

 Horumon To Rinsho 1981 Mar;29(3):305-10 [Jpn]
- Sveger T; Thelin T. Four-year-old children with alphasub 1-antitrypsin deficiency. Clinical follow-up and parental attitudes towards neonatal screening. Acta Paediatr Scand 1981;70(2):171-7
- Szabo B. [Areas of applicability of thin-layer chromatography. Practical mass screening; data on intestinal mucosa lesions]. Padiatr Padol 1983;18(2):105-11 [Ger]
- Szabo L; Somogyi C; Mate M. Experience based on 800,000 newborn screening tests of the Budapest Phenylketonuria Centre. Acta Paediatr Hung 1985;26(2):113-25
- Szeinberg A; Cohen BE. Early blood sampling in neonatal programs for the detection of phenylketonuria. Padiatr Padol 1982;17(2):287-92

- Tableman B; Katzenmeyer M. Infant mental health services: a newborn screener. Prev Hum Serv 1985 Summer;3(4):21-33
- Tada K; Tateda H; Arashima S; Sakai K; Kitagawa T; Aoki K; Suwa S; Kawamura M; Oura T; Takesada M; et al. Follow-up study of a nation-wide neonatal metabolic screening program in Japan. A collaborative study group of neonatal screening for inborn errors of metabolism in Japan. Eur J Pediatr 1984 Aug;142(3):204-7
- Tada K; Tateda H; Arashima S; Sakai K; Kitagawa T; Aoki K; Suwa S; Kawamura M; Oura T; Takesada M; Kuroda Y; Yamashita F; Matsuda L; Naruse H. Intellectual development in patients with untreated histidinemia. A collaborative study group of neonatal screening for inborn errors of metabolism in Japan. J Pediatr 1982 Oct; 101(4):562-3
- Takagi Y; Kanno S; Yamada M; et al. [Clinical significance of C-reactive protein (CRP) for screening test of infection in high risk neonates. Correlation with acute phase reactants score (APR-Sc)]. Acta Neonatol Jpn 1982;18(2):253-8 [Jpn]
- Takanayagi M. [Thirty cases of histidinemia detected by neonatal screening for inborn errors of metabolism]. Chiba Med J 1983;59(5):319-26 [Jpn]

- Talbot HW; Sumlin AB; Naylor EW; Guthrie R. A neonatal screening test for argininosuccinic acid lyase deficiency and other urea cycle disorders. Pediatrics 1982 Oct;70(4):526-31
- Talbot HW Jr; Naylor EW; Guthrie R. Neonatal urine screening for metabolic disease with auxotrophic strains of Bacillus subtilis. Clin Chim Acta 1982 Mar 12;119(3):345-9
- Tan KL; Loganath A; Roy AC; Goh HH; Karim SM; Ratnam SS. Cord plasma alpha-fetoprotein values and neonatal jaundice. Pediatrics 1984 Dec; 74(6):1065-8
- Taswell HF; Reisner RK. Prevention of Rho hemolytic disease of the newborn: the rosette method-a rapid, sensitive screening test. Mayo Clin Proc 1983 May;58(5):342-3
- Taussig LM; Boat TF; Dayton D; et al. Neonatal screening for cystic fibrosis: Position paper. Pediatrics 1983;72(5):741-5
- Tedesco TA. Infant screening for galactosemia in Florida. J Fla Mcd Assoc 1983 Sep;70(9):783-5
- Tejedor JC; Omenaca F; Garcia Caballero C; Oliver A; Quero J; Peralta A; Morreale G; Escobar del Rey F. [Early detection of congenital hypothyroidism. Study of 31,609 determinations]. An Esp Pediatr 1983 May;18(5):364-70 [Spa]
- Thalhammer O. [Screening for congenital hypothyroidism in Austria]. Klin Padiatr 1981;193(5):375-7 [Gcr]
- Thalhammer O; Scheibenreiter S; Knoll E; et al. [12 years Austrian newborn screening for inborn errors of metabolism]. Klin Padiatr 1980;192(6):589-98 [Ger]
- Thelin T; McNeil TF; Aspegren Jansson E; Sveger T. Psychological consequences of neonatal screening for alpha 1-antitrypsin deficiency. Parental reactions to the first news of their infants' deficiency. Acta Paediatr Scand 1985 Sep;74(5):787-93
- Thelin T; McNeil TF; Aspegren Jansson E; Sveger T. Psychological consequences of neonatal screening for alphasub lantitrypsin deficiency (ATD). Acta Paediatr Scand 1985;74(6):841-7
- Thelin TH; Uemura HS; Hsia YE. Symptomatic screening for inborn errors of metabolism. Hawaii Med J 1980 Jun;39(6):139-43
- Therrell BL. Blood from all newborns [letter]. Pediatrics 1982 Jul; 70(1):156-7
- Therrell BL Jr; Brown LO; Dziuk PE; Peter WP Jr. The Texas Newborn Screening Program. Tex Med 1983 Feb; 79(2):44-6

- Tjandra H; Sarwono E. Single step gastric aspirate shake test as a screening procedure for predicting the risk of neonatal respiratory distress. Paediatr Indones 1981 Mar-Apr;21(3-4):69-80
- Topley JM; Rogers DW; Stevens MC; Serjeant GR. Acute splenic sequestration and hypersplenism in the first five years in homozygous sickle cell disease. Arch Dis Child 1981 Oct;56(10):765-9
- Torresani T; Qing Q; Illig R. Thyrotropin enzyme-immunoassay in dried blood spots: A spectrophotometric method for neonatal thyroid screening. J Clin Chem Clin Biochem 1986;24(3):199-203
- Townes PL. Newborn screening: a potpourri of policies [editorial]. Am J Public Health 1986 Oct;76(10):1191-2
- Travert G; Duhamel JF. [Systematic neonatal screening for mucoviscidosis using an immunoreactive trypsin blood assay. Evaluation of 80,000 tests]. Arch Fr Pediatr 1983 Apr;40(4):295-8 [Fre]
- Travert G; Lemonnier F; Fernandez Y. [The direct measurement of free T4 in the cluate of dried blood usefulness as a confirmatory screening test for neonatal hypothyroidism]. J Biophys Med Nucl 1984;8(4):231-4 [Fre]
- Travert G; Mustin C; Fernandez Y. [Newborn screening of mucoviscidosis. Trypsine assay in blood sample]. J Genet Hum 1980;28(4):141-4 [Fre]
- Tredwell SJ; Bell HM. Efficacy of nconatal hip examination. J Pediatr Orthop 1981;1(1):61-5
- Tsukerman GL. [Simple method of mass screening for phenylketonuria]. Lab Delo 1985;(6):326-7 [Rus]
- Tymstra T. False positive results in screening tests: experiences of parents of children screened for congenital hypothyroidism. Fam Pract 1986 Jun;3(2):92-6

U

- Uhrmann S; Marks KH; Maisels MJ; et al. Frequency of transient hypothyroxinaemia in low birthweight infants. Potential pitfall for neonatal screening programmes. Arch Dis Child 1981;56(3):214-17
- Umeda M; Ito Y; Kuroda Y; Enomoto H; Irie M. [The evaluation of the screening kit for cretinism "Daiichi" for TSH assay using filter paper blood samples (author's transl)]. Horumon To Rinsho 1981 Jan;29(1): 115-20 [Jpn]
- Umehashi H; Hayashida T; Otomo S; Fujimoto S; Matsuda I. [The evaluation of measurement of TSH in blood spot on dried filter paper using enzymeimmunoassay]. Horumon To Rinsho 1983 Fcb;31(2):169-72 [Jpn]

- Update on newborn screening. Iowa Mcd 1986 Mar;76(3):139-40
- Urqhuart GE; Carson HG. Laboratory policy for rubella screening [letter].

 Lancet 1982 Nov 27;2(8309):1226

v_____

- Vaca G; Sanchez-Corona J; Olivares N; Medina C; Ibarra B; Cantu JM. On the screening for inborn errors of galactose metabolism. Ann Genet (Paris) 1983;26(3):191-2
- Valman HB. The first year of life: Routine examination of the newborn. Br Med J 1980;280(6207):91-4
- Van Biervliet JP; Rosseneu M. [Screening for dyslipidemia in the newborn infant]. Rev Med Liege 1984 May 15;39(10):469-73 [Fre]
- Van Biervliet JP; Vercaemst R; De Keersgieter W; Vinaimont N; Caster H; Rosseneu M. Evolution of lipoprotein patterns in newborns. Acta Paediatr Scand 1980 Sep;69(5):593-6
- Van Biervliet JP; Vinaimont N; Caster H; Rosseneu M; Belpaire F. A screening procedure for dyslipoproteinemia in the newborn. Apoprotein quantitation on dried blood spots. Clin Chim Acta 1982 Apr 8;120(2): 191-200
- van Gennip AH. [Screening for disorders of purine and pyrimidine metabolism]. Tijdschr Kindergeneeskd 1982 Feb;50(1):30-1 [Dut]
- Varma SK; Cannon C; Varma K; Dooley L. Measurement of reverse triiodothyronine for the newborn screening of thyroid disorders. Indian J Pediatr 1982 Mar-Apr;49(397):185-8
- Velati C; Sampietro M; Sciariada L; Allievi E; Mosconi L; Cappellini MD; Fiorelli G. Neonatal screening for Hb Bart's in Italian subjects of heterogeneous regional origin born in Lombardy. Haematologica (Pavia) 1983 Jan-Feb;68(1):20-9
- Verma UL; Archbald F; Tejani NA; Handwerker SM. Cerebral function monitor in the necnate. I: Normal patterns. Dev Mcd Child Neurol 1984 Apr;26(2):154-61
- Virtanen M; Perheentupa J; Maenpaa J; Pitkanen L; Pikkarainen J. Finnish national screening for hypothyroidism. Few false positives, early therapy. Eur J Pediatr 1984 Nov;143(1):2-5
- Visser HKA. [General screening for congenital hypothyroidism of all neonates in the Netherlands]. Ned Tijdschr Geneeskd 1980;124(13):460-1 [Dut]

W

- Wada Y; Nakajima H; Irie M; Hirayama M; Suwa S. Newborn mass screening in Japan--1984. Jinrui Idengaku Zasshi 1984 Sep;29(3):277-86
- Wadlington, Walter; Waltz, Jon R.; Dworkin, Roger B. Technology and values in transition: the role of law. In: <their> Cases and Materials on Law and Medicine. Mineola, N.Y.: Foundation Press, 1980. pp. 688-1032
- Wadman SK; de Bree PK; Duran M; de Jonge HF. Detection of inherited adenylosuccinase deficiency by two dimensional thin layer chromatography of urinary imidazoles. Adv Exp Med Biol 1986;195 Pt A: 21-5
- Walfish PG; Gera E; Wood MM. Problems in screening for congenital hypothyroidism using thyroxine assays from small dried blood discs. J Nucl Med 1981 Sep;22(9):818-23
- Walker V; Clayton BE; Ersser RS; et al. Hyperphenylalaninacmia of various types among three-quarters of a million neonates tested in a screening programme. Arch Dis Child 1981;56(10):759-64
- Wallace AM; Beastall GH; Cook B; Currie AJ; Ross AM; Kennedy R; Girdwood RW. Neonatal screening for congenital adrenal hyperplasia: a programme based on a novel direct radioimmunoassay for 17-hydroxyprogesterone in blood spots. J Endocrinol 1986 Feb;108(2):299-308
- Walterspiel JN; Rutledge JC; Bartlett BL. Fatal acute splenic sequestration at 4 months of age. Pediatrics 1984 Apr;73(4):507-8
- Wang TR. Application of a thin-layer chromatography microtechnique for the screening of aminoacidurias in 3,165 Chinese newborns. Taiwan I Hsueh Hui Tsa Chih 1984 Sep;83(9):869-78
- Warren NS; Carter TP; Humbert JR; Rowley PT. Newborn screening for hemoglobinopathies in New York State: experience of physicians and parents of affected children. J Pediatr 1982 Mar;100(3):373-7
- Warwick WJ. Prognosis for survival with cystic fibrosis: The effects of early diagnosis and cystic fibrosis center care. Acta Paediatr Scand 1982;71(Suppl.301):27-31
- Warwick WJ; Huang NN; Waring WW; Cherian AG; Brown I; Stejskal-Lorenz E; Yeung WH; Duhon G; Hill JG; Strominger D. Evaluation of a cystic fibrosis screening system incorporating a miniature sweat stimulator and disposable chloride sensor. Clin Chem 1986 May;32(5): 850-3
- Watanabe T; Kuroda Y; Naito E; Ito M; Takeda E; Toshima K; Miyao M; Tomita T; Furukawa S: [A case of homocystinuria missed by the newborn screening]. No To Hattatsu 1985 May;17(3):265-8 [Jpn]

- Webb KC; Krishnan V; Katzman G. Newborn hearing screening using the Crib-O-Gram. Arch Otolaryngol Head Neck Surg 1986 Apr;112(4):420-2
- Weitzel D; Peters H; Humburg C. [Sonographic screening of newborn infants]. Ultraschall Med 1984 Dec;5(6):277-80 [Ger]
- Wilchen B; Brown A; Raby J; et al. Newborn screening for metabolic disorders in Australia and New Zealand: Results for 1983. Med J Aust 1985;143(4):159-60
- Wilchen B; Brown AR; Urwin R; Brown DA. Cystic fibrosis screening by dried blood spot trypsin assay: results in 75,000 newborn infants. J Pediatr 1983 Mar;102(3):383-7
- Wilchen B; Chalmers G. Reduced morbidity in patients with cystic fibrosis detected by neonatal screening. Lancet 1985 Dec 14;2(8468):1319-21
- Wilcken B; Towns SJ; Mellis CM. Diagnostic delay in cystic fibrosis: lessons from newborn screening. Arch Dis Child 1983 Nov;58(11):863-6
- Williams GH; Butts FM. Automated neonatal hearing test in a community hospital. South Med J 1982 Jun;75(6):701-3
- Williams JK. Genetic counseling in pediatric nursing care. Pediatr Nurs 1986 Jul-Aug;12(4):287-90
- Wilson BE; Netzloff ML. Congenital hypothyroidism and transient thyrotropin excess: Differential diagnosis of abnormal newborn thyroid screening. Ann Clin Lab Sci 1982;12(3):223-33
- Wilson DA. Ultrasound screening for abdominal masses in the neonatal period. Am J Dis Child 1982 Feb;136(2):147-51
- Wilson DP; Coldwell JG. Newborn hypothyroid screening. The private sector.

 Am J Dis Child 1985;139(7):662-3
- Wolf B; Heard GS; Jefferson LG; et al. Nconatal screening for biotinidase deficiency: An update. J Inherit Metab Dis 1986; 9(Suppl.2):303-6
- Wolf B; Heard GS; Jefferson LG; Proud VK; Nance WE; Weissbecker KA. Clinical findings in four children with biotinidase deficiency detected through a statewide neonatal screening program. N Engl J Med 1985 Jul 4;313(1):16-9
- Wolff OH; Smith I; Carson N; et al. Routine neonatal screening for phenylketonuria in the United Kingdom 1964-78. Br Med J 1981; 282(6277):1680-4
- Wong HB. Diagnosis of inborn errors of metabolism. J Singapore Paediatr Soc 1986;28(1-2):45-53

- Wong HB; Kham Sk. Screening for hacmoglobinopathics/thalassaemia utilising hacmatologic indices. J Singapore Paediatr Soc 1984;26(3-4):170-5
- Wyatt PR. Issues surrounding genetic screening programs [letter]. Am J Public Health 1981 Dec;71(12):1411
- Wyatt PR. Should screening programs be perfect? [letter] Pediatrics 1983 Feb;71(2):301

-			
•			
v			
•			

- Yamaguchi S; Koda N; Eto Y; Aoki K. Quick screening and diagnosis of organic acidemia by NMR urinalysis. J Pediatr 1985 Apr;106(4):620-2
- Yamamoto K. [Hearing screening for neonates by Crib-O-Gram]. Nippon Jibiinkoka Gakkai Kaiho 1985 Sep;88(9):1132-40 [Jpn]
- Yasuda N. A surveillance system for metabolic disorders in Japan. Hum Hered 1984;34(5):321-7
- Yokoya S; Tokuhiro E; Suwa S; Maesaka H. Measurement of the skin urocanic acid content in normal and histidinemic infants. Eur J Pediatr 1983;140(4):330-2

Z_____

- Zabransky S. [Hypothyroidism screening in newborn infants]. ZFA (Stuttgart) 1980 May 20;56(14):969-73 [Gcr]
- Zammarchi E; Bardini MR; Donati MA; Savelli A. [Results of screening the urine of 26,014 newborn infants for amino acid disorders]. Riv Neurobiol 1981 Jul-Dec;27(34):681-6 [Ita]
- Zammarchi E; Bardini MR; Donati MA; Savelli A. [Screening for amino acids disorders in newborn infants in the city of Florence and neighboring towns from 1973 to 1980. Organizational model]. Riv Neurobiol 1981 Jul-Dec;27(34):677-80 [Ita]
- Zezulakova J; Mala I; Skerikova Z. [System of early screening and rehabilitation of endangered neonates in Prague 4]. Cesk Pediatr 1984;39(10):598-602 [Czc]
- Zollner H; Klinke D. [Laboratory diagnosis possibilities for the early detection and confirmation of cystic fibrosis (mucoviscidosis).

 Overview and personal studies]. Z Med Lab Diagn 1983;24(2):101-7
 [Ger]
- Zucchelli GC; Piro MA; Giannessi D; Malvano R. Radioimmunoassay of trypsin-like immunoreactivity in dried blood spots. J Nucl Med Allied Sci 1982;26(1):35-9